PROF. MATTHEW TOM HARRISON (Orcid ID : 0000-0001-7425-452X) DR. ANNETTE COWIE (Orcid ID : 0000-0002-3858-959X)

Article type : Invited Review

Corresponding Author Email ID: Matthew.Harrison@utas.edu.au

Invited Review

Carbon myopia: the urgent need for integrated social, economic and environmental action in the livestock sector

Matthew Tom Harrison^A*, Brendan Richard Cullen^B, Dianne Elizabeth Mayberry^C, Annette Louise Cowie^E, Franco Bilotto^A, Warwick Brabazon Badgery^D, Ke Liu^G, Thomas Davison^F, Karen Michelle Christie^A, Albert Muleke^A and Richard John Eckard^B

^A Tasmanian Institute of Agriculture, University of Tasmania, 16-20 Mooreville Rd, Burnie, TAS, 7320, Australia

^B Faculty of Veterinary and Agricultural Sciences, University of Melbourne, Parkville, Victoria 3010, Australia

^c CSIRO Agriculture and Food, 306 Carmody Rd, St Lucia, Queensland 4067, Australia

^D NSW Department of Primary Industries, Agricultural Research Institute, Orange, NSW 2800, Australia

^ENSW Department of Primary Industries / University of New England, Armidale, NSW, 2351, Australia

^F Livestock Productivity Partnership, University of New England, Armidale, Australia

^G Hubei Collaborative Innovation Centre for Grain Industry/School of Agriculture, Yangtze University, Jingzhou, China

* Corresponding author

Submitted 30 March 2021; revised version submitted 19 July 2021.

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please cite this article as doi: 10.1111/GCB.15816

Keywords

Emissions intensity, carbon neutral, socio-economic, policy, climate change, adaptation, multidisciplinary, maladaptation, sustainable development goals, carbon dioxide removal (CDR)

Abstract

Livestock have long been integral to food production systems, often not by choice but by need. While our knowledge of livestock greenhouse gas emissions (GHG) mitigation has evolved, the prevailing focus has been - somewhat myopically – on technology applications associated with mitigation.

Here, we (1) examine the global distribution of livestock GHG emissions, (2) explore social, economic and environmental co-benefits and trade-offs associated with mitigation interventions, and (3) critique approaches for quantifying GHG emissions.

This review uncovered many insights. First, while GHG emissions from ruminant livestock are greatest in low and middle-income countries (LMIC; globally, 66% of emissions are produced by Latin America and the Caribbean, East and southeast Asia, and south Asia), the majority of mitigation strategies are designed for developed countries. This serious concern is heightened by the fact that 80% of growth in global meat production over the next decade will occur in LMIC. Second, few studies concurrently assess social, economic and environmental aspects of mitigation. Of the 54 interventions reviewed, only 16 had triple-bottom line benefit with medium-high mitigation potential. Third, while efforts designed to stimulate adoption of strategies allowing both emissions reduction (ER) and carbon sequestration (CS) would achieve the greatest net emissions mitigation, CS measures have greater potential mitigation and co-benefits.

The scientific community must shift attention away from the prevailing myopic lens on carbon, towards more holistic, systems-based, multi-metric approaches that carefully consider the *raison d'être* for livestock systems. Consequential life-cycle assessments and systems-aligned 'socio-economic planetary boundaries' offer useful starting points that may uncover leverage points and cross-scale emergent properties. Derivation of harmonised, globally-reconciled sustainability metrics requires iterative dialogue between stakeholders at all levels. Greater emphasis on the simultaneous characterisation of multiple sustainability dimensions would help avoid situations where progress made in one area causes maladaptive outcomes in other areas.

1. Introduction

Contemporary scientific literature is laden with calls to reduce livestock greenhouse gas (GHG) emissions to limit dangerous climate change (Alcock et al., 2015; Harrison et al., 2014a; Harrison et al., 2016b; Harrison et al., 2014b; Ho et al., 2014; Rawnsley et al., 2018; Schellnhuber et al., 2006). On the other hand, *ceteris paribus* demographic trends suggest that agricultural productivity must increase substantially to meet the demands of a burgeoning and increasingly affluent global population (Struik et al., 2014). These apparent contradictions occur against a background of increasing frequencies of extreme climatic events that impact feed supply, water quality and resources, animal health, infrastructure, profitability and smallholder livelihoods. Together, these factors make the sustainable provision of safe and affordable livestock products increasingly difficult (Bell et al., 2015; Bell et al., 2013; Chang-Fung-Martel et al., 2017; Godde et al., 2021; Harrison et al., 2017b; Harrison et al., 2016a).

Policies designed to achieve net zero or net negative emissions to stabilise the global climate by the end of the 21st century (IPCC, 2014; Schleussner et al., 2016) may limit the potential for increased food production as well as the continued support of rural livelihoods in the decades ahead (Frank et al., 2017; Hasegawa et al., 2015; Smith et al., 2013). Large-scale afforestation and the use of biomass for energy production (Frank et al., 2017; Kreidenweis et al., 2016; Popp et al., 2017) together with population growth will further exacerbate competition for land and food supply.

There are also questions about the applicability of existing GHG emissions mitigation strategies across livestock farming systems (e.g. intensive industrialised livestock production vs extensive rangeland grazing systems), regions and demographics (FAO, 2019). This is particularly important in low- and middle-income countries (LMIC), where the consumption of animal-derived products is growing rapidly: between 1970 and 2012, demand for animal products increased four-fold (FAO, 2019). Collectively, these issues elicit polarising questions of how mitigation of GHG from agricultural systems can take place while productivity is sustainably increased to meet the needs of a growing global population.

The pipeline of sustainable intensification solutions for livestock production systems ranges from theoretical and anecdotal, to proposed, adopted and proven. Sustainable intensification is an intervention that proposes greater agricultural output while keeping the ecological footprint as small as possible (Struik et al., 2014), though multi-disciplinary assessments of sustainable intensification strategies are uncommon. In this article, we examine intervention options through a lens of *emissions reduction* (ER) or *carbon sequestration* (CS): the former reduce *status quo* emissions (e.g. vaccines to reduce enteric methane (CH₄)), while CS options counterbalance GHG emissions per unit area by withdrawing CO₂ from the atmosphere (e.g. planting trees on farm to compensate for livestock CH₄ emissions). ER technologies and practices can only reduce absolute emissions to zero, while CS interventions transfer carbon from the atmosphere into organic forms. Collectively, CS options can exceed the absolute GHG emissions associated with a livestock product, such that net GHG emissions are negative. The combination of ER and CS technologies provides significant potential to maintain or increase livestock productivity while reducing net GHG emissions, which together would reduce emissions intensity (carbon footprint) of livestock products.

While past work has explored ER and CS practices *en masse*, most studies focus either on knowledge, technologies or mitigation potential in isolation (e.g., the effects of a feed additive on enteric CH₄ fermentation) using a limited number of metrics (e.g., GHG emissions and livestock production). Few studies explore the co-benefits and trade-offs associated with GHG mitigation on environmental, economic and social factors (Mayberry et al., 2019), such as nutrient leaching, soil

erosion, biodiversity, profitability, social licence to operate and other socio-cultural factors. These sustainability dimensions underpin the United Nations Sustainable Development Goals (SDGs), which aim to end individual priorities and encourage development within a social, economic and environmental nexus (Focus, 2021).

Here, our objectives were to (1) examine the global distribution of livestock production and highlight regions responsible for the greatest livestock GHG emissions, (2) explore co-benefits and trade-offs associated with economic, environmental and social implications of GHG emissions mitigation, (3) critique current approaches used to quantify and report livestock GHG emissions and (4), highlight opportunities for future GHG emissions mitigation research.

In the following sections, we first outline global livestock production and projected changes in GHG emissions trends, including differences expected between developed countries and LMIC. We then summarise a range of economic, environmental and social co-benefits and trade-offs associated with ER and CS interventions, giving special attention to co-benefits associated with CS in soils and vegetation. The case for cross-sector, multi-metric, multi-scale holistic systems-based emissions mitigation foci is then described, including harmonised solutions for a way forward. We end briefly on research and development (R&D) funding associated with GHG emissions mitigation, including variability in funding cycles.

2. Why are emissions intensities of livestock products generally higher than those of other foods?

Livestock production systems have been criticised for many reasons in the literature, including their contribution to the loss of terrestrial biodiversity (Steinfeld et al., 2006b), their disproportionately low contribution of dietary energy and protein (18% and 39% respectively) relative to their land-use footprint (pasture accounts for 68% of agricultural land, plus one third of cropland is used for animals (Alexander et al., 2015)), competing use of crop production for feed, impact on nutrient surpluses (Menzi et al., 2010; Stehfest et al., 2013), contribution to deforestation (Smith et al., 2013) and detrimental impacts on ecological resilience (Bouwman et al., 2002). A key focus in recent times has been the contribution of livestock to global GHG emissions and climate change (Rose and Lee, 2008; Schellnhuber et al., 2006).

Globally, livestock are responsible for around 14.5% of total anthropogenic GHG (Gerber et al., 2013) and 70% of total emissions from agriculture, forestry and other land use (AFOLU) (Caro et al., 2018). Emissions attributed to livestock systems include carbon dioxide (CO_2) from land use change and burning of fuels, nitrous oxides (N_2O) from fertilisers, and methane (CH_4) from manure and enteric fermentation (FAO, 2021a). Beef and dairy cattle are responsible for the majority of emissions, contributing 64-78% of emissions from the livestock sector (Fig. 1). The majority of cattle emissions are derived from land use change (45%) and enteric fermentation (39%), with smaller contributions from manure storage, meat processing and transportation.

The emissions intensity of livestock products is typically greater than other foods due to low land use efficiency, production of enteric CH_4 (if ruminant based, noting that CH_4 contributes 28 times the global warming of CO_2^{-1}) and emissions associated with land conversion for grazing and/or

¹ Noting caveats associated with global warming potentials explored later in this article

production of animal feed (Ritchie and Roser, 2020). Emissions from ruminants are higher than those from monogastrics because ruminants generally consume lower quality feeds (Herrero et al., 2013b) and have lower feed conversion efficiencies (e.g. the quantity of feed required to produce one kilogram of beef or lamb/mutton is around 25 kg or 15 kg respectively, whereas that for pork and poultry is 6.4 kg and 3.3 kg respectively (Alexander et al., 2016; Ritchie and Roser, 2020). Although emissions intensities for agricultural products vary widely across commodities, region and production systems, averaged globally and across the supply chain, beef, lamb/mutton, pig meat, poultry meat, dairy milk emissions intensities are 50, 20, 7.6, 5.7 and 3.2 kg CO₂-e/kg food product respectively. These contrast with the much lower emissions intensities of plant-based products including wheat, maize, pulses and root vegetables, which have emissions intensities of less than 0.8 kg CO₂-e/kg product (Poore and Nemecek, 2018).

Does this mean that we should cease all livestock production systems? Perhaps not, at least until we have more carefully explored the *raison d'être* of livestock systems, including biophysical, economic, social and cultural reasons, and considering the fact that global variability in emissions intensities of ruminants is greater than most other foods. For example, the production of 100 grams of beef protein is responsible around 25 kg CO₂-e, but this number hides enormous variability, with 10th and 90th percentiles ranging from 9 kg to 105 kg CO₂-e respectively (Poore and Nemecek, 2018). This variability - even among producers in similar geographic regions - implies substantial potential to mitigate environmental impacts and enhance the productivity of the food system (Poore and Nemecek, 2018).

3. Why do we need livestock?

If livestock production systems 'contribute to environmental problems on a massive scale' (FAO, 2006a), why are they still practiced today? For many rural communities, livestock satisfy a variety of societal, economic and environmental needs. In addition to the production of food and fibre, livestock provide income, draught power, nutrient recycling through manure, and have important social and cultural roles that need to be considered (Steinfeld et al., 2003). For example, livestock have important societal values, such as status or dowry (bridal gifts), and are a vital part of many religious and cultural ceremonies such as Eid al Adha, the Muslim festival of sacrifice. As well, some religions prefer red meat over white meat, and Islamic and Jewish faiths forbid consumption of pork (Brondz, 2018; Larson, 2000; Nakyinsige et al., 2012; Rohman and Che Man, 2012).

Livestock are a fundamental component of the economies of LMIC, providing nourishment and income for rural communities, especially women and nomadic populations (Kristjanson et al., 2014; Rubin et al., 2010). It follows that livestock are a crucial asset and safety net for a large portion of the world. In many regions, livestock comprise the main if not only capital reserve of farming households, serving as a strategic and fluid reserve that reduces risk, provides food and non-food products (e.g. wool, hides and skin (Thornton, 2010)), and facilitates income diversification and flexibility in response to changing economic and environmental conditions (Kristjanson et al., 2014; Rubin et al., 2010).

In drier regions where aridity is too high for reliable crop production, livestock are often the only viable option (Thornton, 2010; Thornton and Gerber, 2010). Of the 51 million km² of habitable global land, livestock occupy 40 million km² (Mottet et al., 2017; Ritchie, 2019), including grazing land and arable land used for animal feed production (Ritchie and Roser, 2020). Although short-cycle monogastric production systems such as chickens and pigs can be important for household food

security and immediate cash flow, it is worth stressing that only ruminants are able to convert fibrous material and forages - that would otherwise have little or no alternative use - into valuable products (FAO, 2019). Each year, ruminants convert some 2.7 billion metric tonnes of grass dry matter, of which 65% grows on land unsuitable for crops, into edible human protein (Mottet et al., 2017). In developed countries, there is a negative, near-linear relationship between cost of production and proportion of pasture in the diet (Dillon, 2007), implying that diets comprised by greater proportions of grazed material are conducive to greater profitability.

If a new food production system was proposed to replace or displace livestock on lands unsuited to crops, the new system should also be subject to environmental and GHG emissions audits, with the emissions intensity of the new product compared with that from the preceding livestock system, while appropriately accounting for the multi-functional values of some livestock systems (Weiler et al., 2014). A fundamental flaw of many past studies has been the lack of comparison of emissions associated with livestock products with alternative (but realistic) agricultural products derived from the same environment. Such assessments can be conducted using *consequential life cycle assessments*, and are the focus of a later section in this article.

As explained below, these essential socio-economic, environmental and cultural roles of livestock are becoming increasingly important as the sector grows (Herrero et al., 2013a). The dependence of millions of poor people on livestock suggest that climate change mitigation policies involving livestock must be designed with extreme care (Havlík et al., 2014). Any political incentive intended to reduce livestock supply or demand should first very carefully consider the global distribution of livestock as a function of the biophysical, socio-economic and cultural services they provide, particularly in locations where livestock underpin the necessities of life.

4. How (and where) will global livestock supply, demand and GHG emissions change in future?

Increasing human population and the global trend towards urbanisation will drive increasing consumption of livestock products over the next 20 years (FAO, 2006a, b), particularly in LMIC (Herrero et al., 2016). With rising affluence, the consumption of milk, meat and eggs tends to increase (Fig. 2), contributing to improved nutrient status and health of vulnerable populations (Steinfeld et al., 2003). In contrast, increasing incomes in developed countries are no longer associated with increasing animal product consumption. Indeed, higher incomes in developed countries have led to a steady decline in animal food consumption associated with health concerns such as heart and blood circulation diseases (Rae and Nayga, 2010).

Dietary shifts in the emerging economies of Asia and Latin America have placed unprecedented demand on animal products (Godde et al., 2018). Beef and milk production have more than doubled over the last 40 years, with monogastric production (pigs and poultry) in some areas growing by a factor of five or more (Thornton, 2010). In 2019, global meat production was 340 Mt per annum (>35% poultry meat, 35-40% pig meat, and 22% for beef and buffalo meat combined), with production dominated by Asia (144 Mt), followed by Europe (64 Mt), and North and South America (52 and 46 Mt respectively) (FAOSTAT, 2020). Meat production is expected to rise to 366 Mt by 2029 (FAOSTAT, 2020). Similarly, the 852 Mt of milk produced globally in 2021 is expected to grow at 1.6% per annum to 997 Mt by 2029, faster than many other agricultural commodities (Fig. 3).

Globally, 66% of livestock emissions are produced by Latin America and the Caribbean, East and southeast Asia, and south Asia (FAO, 2021b). Latin America and the Caribbean produce nearly 1.9 Gt CO₂-e/annum associated with beef production and land-use change (deforestation and pasture

expansion). East and Southeast Asia are the second largest emitter (1.6 Gt CO_2 -e/annum), followed by South Asia (1.5 Gt CO_2 -e/annum). North America, Western Europe, East and North Africa each produce around 0.6 Gt CO_2 -e/annum, but these African regions produce less than half the protein of North America and Western Europe (FAO, 2021b). Sub-Saharan Africa produces around 0.4 Gt CO_2 e/annum. Globally, emissions from livestock systems increased by 51% between 1960 and 2010 due to growth in livestock production (FAO, 2021b); emissions growth is projected to continue until at least 2050, *ceteris paribus* (Fig. 4). The majority of emissions growth has historically occurred in Asia, Africa and the Americas, primarily associated with ruminant production (Fig. 1).

Growth in livestock emissions has been influenced by many interacting socio-economic, biophysical, political and environmental factors. Asian and African regions have each increased livestock GHG emissions since 1970, but for different reasons. In China, government policies, growing corporate integration and urbanisation have encouraged meat and milk consumption (Bai et al., 2018). In contrast, livestock production in the low-income regions of Africa has been driven by higher animal numbers associated with expansion of grazing areas with low agroecological potential, coupled with low market access, reduced government investment and high labour costs (Godde et al., 2018). For these reasons, higher livestock production in Africa has been accompanied by higher GHG emission intensities for some livestock products (Fig. 5), increasing the emissions from the livestock sector. In contrast, GHG emissions from livestock in developed countries have decreased by 23% from 1960 to 2010 (Caro et al., 2014). Many countries in Europe and Oceania have transitioned to towards extensification (similar production per unit area from a greater total area) and deintensification (lower per unit area production) and to ethically- and environmentally-friendly livestock products, decreasing either gross GHG (Fig. 4) and/or emission intensities (Fig. 5). Such trends underscore an urgent need for significant improvement in livestock production efficiency in LMIC. This challenge is underpinned by the need to sustainably increase animal production without degrading natural capital or harming the global environment (Box 1).

Box 1: One of the greatest challenges of the 21st century

Satisfying the growing demand for animal products while sustaining the natural resource base (soil, water, air and biodiversity) is one of the foremost challenges facing the world today (FAO 2019; 2021a). Future global agriculture will be increasingly driven by trends in the livestock sector, many of which are already apparent (FAO 2006; 2019; 2021a):

- 1. Livestock production will become more prone to pests and infectious zootonic diseases, which may have implications for public health such as the COVID-19 pandemic that devasted global health and economies from 2019;
- 2. Livestock production systems will shift from local multi-purpose activities towards intensive, market-oriented and increasingly integrated supply chains;
- 3. Consumption and competition (domestic, national and international) for shared natural resources will increase;
- Agricultural systems will transition towards large-scale industrial systems situated near urban centres, but also towards sources of feedstuff, whether crops or trade hubs where feed is imported;
- 5. Pig and poultry production will increase in importance relative to ruminant production, tending to drive down emissions intensities from the livestock sector; and
- 6. The majority of growth in meat production in the next 20 years will occur in Asia and the Americas.

Meeting these challenges will require concerted efforts in global and public policy, designed to improve equity, alleviate poverty and improve environmental, public health and food safety. Such efforts must be underpinned by international coordination with strong leadership.

5. Economic, social, environmental and biophysical co-benefits and tradeoffs caused by GHG emissions mitigation interventions

To examine the extent to which previous work has been conducted with multi-disciplinary collaboration, we searched the Web of Science Core Collection database for documents produced between 1945 and 2021, incrementally adding terms to the baseline 'greenhouse gas emissions mitigation AND livestock' search. Results are shown in Fig. 6a.

Adding the term 'product*' reduced the search results from 804 to 648 (where '*' represents a wildcard such that any text containing 'product' will obtain a result, e.g. production, productivity etc.). Our search revealed a lack of studies canvassing two or more disciplines associated with livestock GHG emissions mitigation. Many studies focussed on technological aspects, such as mitigation potential, livestock productivity, and management of land and/or livestock.

Papers including environmental aspects were lacking (reduction in search results from 804 to 348 in Fig. 6a) but perhaps most stark was the dearth of social studies of livestock GHG emissions mitigation, with only 44 relevant documents retrieved. There were even fewer reports of concurrent economic, environmental and social aspects associated with livestock GHG emissions mitigation (Fig. 6a). To further examine the extent to which transdisciplinary studies have focused on livestock emissions mitigation options for LMICs, we repeated the search in Fig. 6a adding the terms 'AND (LMIC OR (low AND Middle AND income) OR "developing countr*")'. Searches including the terms 'economic', 'environmental' or 'social' respectively revealed only 23, 20 and 6 relevant documents from the entire Web of Science database (Fig. 6b), clearly signifying a lack of multidisciplinary livestock mitigation studies for LMIC. The limited number of social studies likely reflects the order in which traditional reductionist research historically occurs, first with GHG emissions mitigation and biophysical/productivity potential, and last with environmental, social and political factors (Fig. 7).

Using the search results shown in Fig. 6, we next reviewed the economic, environmental and social co-benefits and trade-offs associated with ER and CS interventions (Table 1). These were grouped and ranked in ascending order of increasing economic, social and environmental benefit within each of the nine mitigation categories examined. Few interventions had high mitigation potential with concurrent economic, environmental and social benefit. By way of example, feeding nitrate to beef cattle in rangeland environments has been advocated as a pathway by which non-protein nitrogen could be supplemented to cattle in dry seasons to improve productivity (MLA, 2016, 2020). In Australia, feeding nitrate as a supplement to grazing beef cattle is a legislated CS method under the Commonwealth Emissions Reduction Fund (a policy where farmers and landholders earn carbon credit units and financial return for practices that reduce GHG emissions). While nitrate supplementation has GHG emissions mitigation potential in some environments, it can be unprofitable due to limited effects on productivity (Harrison et al., 2016b) and may have animal welfare implications through nitrate poisoning if the feeding regime is inappropriate (Table 1; Callaghan et al., 2014; Callaghan et al., 2021; MLA, 2016).

Of the 48 ER interventions shown in Table 1, 18 had triple bottom-line benefit. All manure management interventions had economic, environmental and social benefit, with most being applicable in LMIC and having high GHG mitigation potential due to their ability to limit both CH₄ and N₂O emissions (through effects on methanogenesis and/or denitrification). However, manure management approaches are generally applicable only to intensive production systems such as dairies, feedlots and piggeries in developed countries, and smallholder confined feeding operations and subsistence farming operations in LMIC, because manure must be accessible and manageable. For the majority of grassland-based systems, it is not feasible to collect livestock faeces and urine because excreta is spread too sparsely. Such areas include the dryland tropics and continental locales of Asia and North America (Petersen et al., 2013; Steinfeld et al., 2006a) as well as rangelands in central Australia, the African savannas, the South American Pampas and the Great Plains of North America.

For pasture and animal management, only those interventions designed to increase the growth rates of young livestock (or production efficiency), reduce age at slaughter or increase fecundity/weaning rates had medium-high mitigation potential. Reducing time to slaughter and/or improved growth rates of young animals benefits farmers (1) economically through greater liveweight production, (2) environmentally through lower residence time of animals on farm which, at broader scales, causes less soil erosion and destruction of natural habitats, and (3) socially through improved animal welfare and lower mortality rates of juvenile and adult animals. However, any intervention that improves growth rates of young animals only has mitigation benefit if overall stocking rates are not subsequently increased, e.g. if animals removed from a farm are not replaced by others. This highlights an important distinction between reductionist mitigation research versus approaches that evaluate livestock emissions as part of a holistic system; only integrated systems approaches capture important trade-offs between mitigation at the animal and farm level, and between the farm and global level. Such trade-offs may include efficiency gains at the animal level that are offset by increased numbers of animals on farm over a set duration. 'Pollution swapping' is another trade-off

that can be captured by systems approaches, for example increased N fertiliser can improve pasture growth and SOC sequestration, but this may result in greater N₂O emissions. Another trade-off is 'leakage', such that emissions mitigation on one farm is offset by increased emissions on another farm (noting that leakage could be temporal, sectoral or geographical). Leakage may include (1) destocking a farm to reduce emissions, leading to increased stocking rates elsewhere to maintain supply, and (2) revegetation or bioenergy cropping on pasture land that leads to deforestation elsewhere to provide pasture. Underpinning many of the animal management options that have medium-high mitigation potential is reduced stocking rate. This is particularly important in LMIC and extensive areas, because reduced animal densities facilitate higher feed per animal, improved growth rates, higher soil CS (Zhang et al., 2015b) and potentially landscape restoration (discussed further below).

Of the 19 CS interventions in Table 1, 6 had triple-bottom line benefit, 5 of which had medium-high potential to reduce emissions. Perennial legumes (e.g. leucaeana and Desmanthus) offer multiple avenues for emissions mitigation for the environments in which they are suited. These include increased soil carbon storage at depth, higher crude protein and mitigation of enteric CH₄ through condensed tannins (Harrison et al., 2015; Radrizzani et al., 2011; Ruis and Blanco-Canqui, 2017; Sándor et al., 2020; Suybeng et al., 2019; Tomkins et al., 2019), though in some cases legumes may increase N_2O emissions as a result of higher deposition of urinary-nitrogen (Harrison et al., 2016b; Harrison et al., 2015). Forage brassicas (e.g. forage rape) also have medium-high mitigation potential due to higher digestibility that can lead to reduced enteric CH₄ per unit dry matter consumed, higher livestock growth rates and potentially increased soil CS (Ruis and Blanco-Canqui, 2017; Sun et al., 2016; Thomson et al., 2016). But dietary composition is also important: if perennial legumes or forage brassicas form only a small part of annual intake, mitigation of enteric CH₄ is likely to be small. This observation highlights the important trade-off between component-level research (e.g. evaluation of CH₄ mitigation per unit intake of the forage per se) versus systems approaches that account for both mitigation per unit intake but also other features that influence GHG mitigation, including pasture botanical composition and animal dietary constituents.

The trade-offs associated with many of the interventions in Table 1 underpin *a clear need for transdisciplinary efforts* in future assessments of mitigation interventions. Many of the interventions have been evaluated only in a theoretical sense and focus entirely on GHG emissions mitigation, thus ignoring important cross-disciplinary factors such as cost, practicality, reversibility, market access, political factors or social license to operate, which collectively and ultimately define adoptability and impact. Such factors limit the feasible mitigation to a value that is much lower (Fig. 7).

The majority of interventions in Table 1 were tested in and *designed for developed countries*. There are fewer mitigation options that have been tested in LMIC. Mitigation options that could be applied in LMIC that have triple bottom-line benefit include improved animal health and animal husbandry leading to greater growth rates and/or younger age at slaughter, feeding of concentrates, planting trees or perennial legumes (e.g. leucaena) and manure management techniques. Only seven of these interventions have medium or high GHG emissions mitigation potential. Similar to developed countries, not all of these options will be applicable in LMIC due to farming systems practices, location, workforce availability, access to markets and agronomic knowledge, cultural aspects and other factors. Cohen et al. (2021) indicate that a greater understanding of the co-impacts (cobenefits and adverse side effects) is needed to realise the potential for sustainably meeting multiple objectives, increasing the efficiency and cost-effectiveness of climate actions (both mitigation and adaptation). They note that the impact of such information is determined by the manner in which it

is framed and communicated, and that framing needs to be context- and application-specific.

The majority of GHG mitigation studies with an economic focus were associated with profitability/cost of a given intervention (e.g. research and development costs, capital investment, costs of feeding, effects of carbon price (Cohn et al., 2014; Wall et al., 2010)), or the opportunity costs of using land and other resources to raise animals considering their alternative use for other food or non-food sources (Garnett, 2009; Wall et al., 2010). We found that more economic studies examine co-benefits or positive effects associated with GHG mitigation, compared with environmental and social studies (Table 2).

In contrast to economic studies, environmental studies were broader in scope and tended to be equally focused on the co-benefits and trade-offs associated with GHG emissions mitigations (black and red text respectively in Table 2). Assessments ranged from environmental implications caused by production of algae for livestock feeds, to the effects of trees or deep-rooted pastures on soil carbon, biodiversity and natural capital, to planting legumes to replace synthetic nitrogen fertilisers. Studies of manure management were also common but less frequently were focused towards LMIC, with the notable exception of the review by Petersen et al. (2013) that discussed a range of solid and liquid manure management options in Sub-Saharan Africa, Southeast Asia, China and Europe. The mitigation potential associated with reduced food loss and waste was high (Bellarby et al., 2013). As there are many livestock production and environmental co-benefits associated with managing soils and vegetation (trees and shrubs), these interventions are given special consideration in the following sections.

As discussed above, there was a *paucity of social studies* compared with other disciplinary areas (Table 2; Fig. 6a). Most social studies were related to animal welfare, including health benefits of feed additives, mortality associated with higher fecundity breeds, maladaptive breeding traits and co-benefits derived from shade and shelter belts. While many studies focused on health or dietary implications associated with GHG mitigation through reduced demand or consumption (e.g. Bajželj et al., 2014; Popp et al., 2010; Tilman and Clark, 2014; Westhoek et al., 2014), few studies focused on participation in carbon markets or motivations/barriers to adoption of mitigation interventions (e.g. Kragt et al., 2017; Torabi et al., 2016). The majority of studies centered on social aspects of carbon farming were conducted in and for developed countries.

Despite considerable variability in the pathways with which co-benefits and trade-offs could be realised with individual mitigation interventions, many share common principles:

- 1. Economic co-benefits:
- <u>Improved liveweight gain and/or animal productivity</u> e.g. feeding of supplements, higher digestibility feed types, improved pasture species, higher fecundity breeds, productivity cobenefits associated with shade or provision of shelter;
- <u>Reduced management/input costs:</u> e.g. reducing dairy cow milking frequencies or using less synthetic fertilisers as a result of planting legumes;
- <u>Income diversification:</u> through access to new markets such as carbon trading, ecosystems services or payments for biodiversity, renewable energy, improvements to water and/or air quality, or the ability to create a new product (e.g. bioenergy crops) or branding ('carbon neutral').

2. Economic trade-offs:

• <u>Cost of implementation</u> e.g. the limited effect of some supplements on liveweight gain (e.g. nitrates) or the small effects of nitrification inhibitors on pasture growth do not compensate for

the cost of the intervention (Callaghan et al., 2021);

- <u>Market variability</u>: reduced prices for animal products, carbon or ecosystems services may result in lower income compared with no intervention;
- Loss of productive arable land due to e.g. planting trees or restoration of natural habitats;

3. Environmental co-benefits:

- <u>Pasture sustainability</u> e.g. improved ground cover, pasture persistence and production resulting from management practices that facilitate spelling or reduced grazing duration of grasslands (such as seasonal confinement feeding or introducing animal genotypes with greater feed-conversion efficiency), reduced need of synthetic fertilisers by planting legumes, converting to permanent pastures as opposed to annual crops;
- <u>Ecosystems services:</u> e.g. improved water quality from large-scale algae production, improved soil carbon from planting deeper-rooted pasture species or spreading biochar, reduced leaching of nutrients into waterways through manure management;
- <u>Reduced need for fossil fuels:</u> preventing/displacing off-farm GHG emissions e.g. planting of bioenergy crops (Goldemberg, 2007), use of renewable energy on farm such as solar panels (photovoltaics), wind turbines or anaerobic digestors for manure (Bardi et al., 2013; Hernandez et al., 2015).

4. Environmental trade-offs:

- <u>Emissions leakage</u> in which emissions no longer occurring in a location occur elsewhere, e.g. emissions caused by production of feed additives or grain used for animal feeds;
- <u>Reduced water quality</u> such as leaching of P and N into waterways, causing eutrophication;
- <u>Reduced soil quality</u> e.g. acidification caused by excessive use of nitrogenous fertilisers or inclusion of legumes in pasture;
- <u>New pasture types may become weeds</u> if unmanaged or allowed to escape;
- <u>Habitat for pests</u> may be engendered if vegetated areas are created (or land spared), providing habitat for vermin, weeds and pests that together may damage both natural and arable areas.

5. Social co-benefits:

- <u>Animal health and welfare</u> through provision of shade and shelter, reduced heat stress, reduced need for labour, improved animal traction in LMIC, improved milking frequencies of dairy cows, improved nutrition (some pasture types and legumes);
- <u>Human health and welfare</u> associated with reduced exposure to heavy metals (manure management), improved water quality from reduced ground water contamination, increased firewood supply associated with afforestation in LMIC, social relationships in LMIC and improved health of rural communities with the addition of animal protein to human diets;
- <u>Enhanced resilience of rural communities</u> through additional expenditure on infrastructure such as fencing to manage over-grazing by livestock and feral herbivores, carbon trading, which provides a consistent source of income during drought, enabling early destocking to minimise risk of land degradation and hasten pasture recovery post-drought (Cowie et al. 2019).

• Social trado

Social trade-offs:

- <u>Animal health and welfare</u> associated with some feed additives (e.g. potentially carcinogenic effects of algal feeds (Abbott et al., 2020) and tannins (Jerónimo et al., 2016) increased mortalities associated with higher fecundity breeds, maladaptive breeding traits);
- <u>Adverse societal impacts:</u> caused by fuel/livestock feed competition, competition for land;

vegetation management at the expense of food security in the absence of adequate governance; livelihoods impacted where livestock are the main source of income, especially in some LMIC, and pollution swapping (e.g. where livestock GHG emissions mitigation facilitates greater use of fossil fuels elsewhere (Smith et al., 2007)).

Carbon sequestration opportunities provide a pathway for greater emissions mitigation than the potential associated with ER. As detailed further below, two promising emissions CS pathways include CS in soils and in vegetation: these options provide medium-high mitigation potential (Table 1) with ability to offset enteric CH₄ (Doran-Browne et al., 2016; Henderson et al., 2015; Leahy et al., 2019; Smith et al., 2020), are widely applicable in developed countries and LMIC, and offer multiple co-benefits.

6. Mitigation potential and co-benefits associated with carbon sequestration in soils

Grazing lands (savannas, grasslands, prairies, steppe, and shrublands) cover 45% of the earth's land surface (excluding Antarctica) (Ritchie, 2019), such that modest increases in soil organic carbon (SOC) could offset GHG emissions from livestock in grazing lands (Derner and Schuman, 2007; Zhang et al., 2015a). Recent work has suggested that soil carbon represents 25% of the potential of natural climate solutions, of which 40% is protection of existing soil carbon, and 60% is rebuilding of depleted stocks (Bossio et al., 2020). Co-benefits associated with increased soil C in grazing systems include improved plant available water holding capacity and increased nitrogen supply through mineralisation (Wander and Nissen 2004), which may buffer the impact of climate variability and change.

There are several grazing and land management options that can increase SOC (Table 1 and 2). Grazing practices that improve SOC include those that alter pasture species composition, forage consumption and ground cover, leading to greater nutrient, organic matter or water ingress into pastures (Conant and Paustian, 2002; Lal, 2004). Land management practices that can increase SOC include (1) moving from annual cropping to perennial pastures (Badgery et al., 2014), (2) increasing soil fertility, (3) improving grazing (in some cases reducing stocking rate) and (4) planting trees or deep-rooted perennials. Altering stocking rate to maximise forage production could result in a global sequestration of up to 148 Tg CO₂/annum in grazing lands (Henderson et al., 2015; Herrero et al., 2016), while sowing legumes could sequester an additional 203 Tg CO₂/annum (despite being applicable over a much smaller area; Henderson et al., 2015). However, increased N₂O emissions from legumes can offset SOC sequestration benefits (Henderson et al. 2015), again suggesting a need to holistically quantify changes in all of the main GHG emissions (CO₂, N₂O, CH₄) whenever an intervention is made.

Because there are many interactions that govern soil carbon flux, grazing often has variable and inconsistent effects on SOC. Grazing perturbs above- and below-ground net primary production, shifting grassland composition (Derner et al., 2006), altering nutrient cycling, groundcover, soil temperature, moisture and respiration. Shifting from heavy to light grazing utilisation may increase SOC (Byrnes et al., 2018), with SOC under light grazing often being higher than that with no grazing (Jiang et al., 2020; Liu et al., 2012; Orgill et al., 2012).

More than 80% of the potential increase in SOC is in LMIC, where pasture production would be expected to increase following periodic de-stocking, resulting in a lower annual average stocking rate (Henderson et al., 2015). East and South Africa in particular have large potential for increased SOC sequestration (0.1-3.1 Mg/ha.annum) and there are reports of up to 90% decline in SOC in the heavily grazed highlands of South Africa (Dlamini et al., 2014). Despite this, very little work has been conducted on the potential of practices or incentives to encourage SOC sequestration in East and South Africa (Tessema et al., 2020).

It is important to note that there are limits to the maximum SOC stock in any soil. If SOC is already close to its maximum, further increases to SOC will be unlikely. In these cases, effort would be better spent in preventing decline of SOC (Badgery et al., 2020; Sanderson et al., 2020) through implementation of the approaches outlined above. Lack of continuing nutrient input (e.g. fertilisation), drought exposure, or other changes relative to historical land management may also enhance the rate of organic carbon conversion to CO_2 (either through leaching or respiration), causing soil carbon to decline.

There are substantial opportunities to increase SOC on degraded grazing lands, because the most degraded soils often have the highest capacity to store additional SOC. This is most evident in LMIC such as East Africa and southwest China where stocking rates are often excessive, impacting on production and causing degradation (Lohbeck et al., 2018; Michalk et al., 2019; Pang et al., 2019). Of the total global potential in SOC sequestration in grazing lands, the largest potential is in Central/South America (26.7 Tg CO₂/annum), Sub-Saharan Africa (24.3 Tg CO₂/annum), Oceania (15.6 Tg CO₂/annum) and East/Southeast Asia (13.7 Tg CO₂/annum) (Henderson et al. 2015). While there is moderate variation in regional sequestration rates (0.13 to 0.32 Mg CO_2 /ha.annum, coefficient of variation = 26%), most of the difference in total carbon storage potential between continents was due to variations in land area (coefficient of variation = 72%). Henderson et al. (2015) also found a relationship between agroecological zone (temperate, humid and arid) and sequestration potential. At the global average level, humid rangelands have the highest C sequestration rates per hectare, followed by arid and temperate rangelands. However, sequestration rates differ to total sequestration potential. Arid areas account for just over half of the total global soil C sequestration potential due to their dominant share of the total rangeland area. At the global level, the largest soil C sequestration potentials in lie in Central and South America, Sub-Saharan Africa and Oceania due to the large areas of rangelands in the humid and arid agroecological zones within these continents. Similarly, the low per hectare potentials in Central Asia, Eastern Europe and Russia, East and Southeast Asia reflect the higher proportions of temperate rangelands in these regions (Henderson et al., 2015).

7. Mitigation potential and co-benefits associated with carbon sequestration in vegetation

Similar to the CS opportunities afforded by soil carbon, the global CS potential in vegetation is large (Domke et al., 2020). The CS potential associated with reforestation (planting trees where there once was forest) and afforestation (planting trees on previously unforested land) is respectively around 1.5–10.1 Gt CO₂-e/annum (Smith et al., 2020) and 1.5 Gt CO₂-e/annum (Nilsson and Schopfhauser, 1995). The majority of this global carbon sink potential is in grasslands (Scurlock and

Hall, 1998).

Forestry (trees only), agroforestry (integrated agriculture and trees) and silvopastoral systems (integrated livestock grazing and trees) are recognised carbon offset activities in some carbon marketing schemes, creating opportunities for landholders in LMIC to generate income through CS associated with vegetation management. Doran-Browne et al. (2018) showed that planting trees on 20% on a high-density livestock farm in Australia achieved carbon neutrality. However, after 30 years, CS in the trees had slowed considerably. This result highlights the contrasting transient nature of CS relative to the continuing emissions of livestock (assuming stocking rates remain unchanged), suggesting a need for interventions that are not limited by saturation once carbon stored in soils and vegetation nears its potential. The sequestration ceiling could be avoided by harvesting forests for use in long term products, harvesting biomass for bioenergy, and producing and applying biochar to soils.

Although agroforestry practices in LMIC are more widespread than in developed countries (Ramachandran Nair et al., 2010), carbon market opportunities in LMIC associated with agroforestry are lower, hampered by insecure land tenure and high transaction costs that are prohibitive for smallholders in Africa and Indonesia for example (Cacho et al., 2005; Jindal et al., 2008). Widespread participation in carbon markets in the foreseeable future is unlikely without accompanying development of policy instruments to reward the rural poor for other environmental services such as watershed protection and biodiversity enhancement (Garrity, 2004; Lee et al., 2016). This type of financial reward may help recoup landholder costs incurred in protecting and enhancing environmental services.

Agroforestry provides a range of ecosystems services. Carbon sequestration in soils and vegetation could perhaps be more appropriately viewed as a co-benefit from improving livestock productivity and ecosystems services, rather than a primary objective for either managing land ecosystems (Henderson et al., 2015; Herrero et al., 2016) or for deriving income from CS *per se* (Alcock et al., 2015; Harrison et al., 2016b; Ho et al., 2014). Planting trees can improve livestock production through shade and shelter that regulates the microclimate (Deniz et al., 2019) which potentially improves liveweight gain and survival (Gregory, 1995; Montagnini et al., 2013), while fodder trees and shrubs can be a valuable source of feed in dry seasons (Franzel et al., 2014; Vandermeulen et al., 2018). In dry environments, hydraulic redistribution (Bayala and Prieto, 2020; Bogie et al., 2018) could contribute towards enhanced growth of pasture under some tree species.

In the low productivity rangelands of developed countries, income from "carbon farming" (e.g. avoided deforestation, afforestation and forest regeneration) can enhance the resilience of farming systems and rural communities, facilitating expenditure on infrastructure such as fencing to manage over-grazing by livestock and feral herbivores, providing income during periods of drought, enabling early destocking to minimise risk of land degradation and hasten pasture recovery post-drought (Cockfield et al., 2019; Cowie et al., 2019). While commercial forestry provides a significant avenue for CS, establishment of industrial plantations in LMIC can displace transhumant pastoralism (Ramprasad et al., 2020) and exacerbating poverty in local communities (Andersson et al., 2016). Commercial forestry companies typically manage multiple stands of different ages to provide a constant supply of products and income. Sawlog rotations are around 25-80 years, pulp rotations 10-15 years, and coppiced biomass rotations are 3-5 years, depending on climate. As income is received

after harvest, delayed returns present a significant cash flow issue for smallholders who own a limited number of stands. However, landholders who plant woodlots usually do so as investments to supplement income from agricultural enterprises. Indeed, Nigussie et al. (2021) found that integration of short-rotation acacia plantations in diversified farming enterprises enhanced livelihoods of some (but not all) smallholders in Ethiopia.

The review comment is applicable to LMIC where international forestry companies develop plantations on leased community land or public land, displacing informal grazing.

Silvopastoral systems provide a range of environmental co-benefits and livestock co-production benefits. Leguminous trees in silvopastoral systems can enhance nutrient cycling and improve soil fertility (Lira Junior et al., 2020; Martínez et al., 2014) and globally have significant potential to sequester carbon (see previous section). Moderate tree cover can increase infiltration and reduce runoff in drylands (Cerdà and Rodrigo-Comino, 2020). Riparian corridors may improve water quality through stabilisation of stream banks and reduction of runoff to waterways (Abernethy and Rutherford, 1999; Hansen et al., 2010). However, pasture reforestation can sometimes decrease soil carbon (Guo et al., 2008) and in some cases, silvopastoral systems can have lower SOC than open pastures (Douglas et al., 2020). The interaction between tree species, soil type and climate has a significant influence on SOC dynamics, and should be considered in GHG accounting for agroforestry (Douglas et al., 2020).

Trees can also enhance the biodiversity of grazing systems (Table 2), particularly where native species are planted. Scattered trees in paddocks or fields and living fences play a key role in ecosystem function and biodiversity conservation in highly modified landscapes (Manning et al., 2006). Plantings that include understorey trees and shrubs increase structural and species diversity, improving habitat value for fauna including pollinators and biological control agents. When strategically located relative to remnant vegetation, reforestation with native species can enhance landscape connectivity, creating biological corridors that may support adaptation, facilitating movement of species in a changing climate (Beier et al., 2012). Trees on farms can improve biodiversity and ecosystem functions indirectly by reducing pressure on natural forests from fuelwood extraction (Table 2).

While there are synergies between reforestation and livestock production, there are trade-offs with agricultural production as tree basal area increases (Table 2; Cerdà and Rodrigo-Comino, 2020). There are also trade-offs between environmental co-benefits: CS is often higher in tree monocultures, while biodiversity value and resilience are greater in mixed species plantings that include shrubs (Paul et al., 2016).

8. The need for integrated, cross-sector, multi-metric, multi-scale and holistic systems-based emissions mitigation foci

While assessment of the *technical* potential for mitigation is a useful first step towards quantifying the *feasible* mitigation associated with an intervention, future studies must go broader to examine the co-benefits (synergies) and trade-offs (antagonisms, both unavoidable and unexpected) that exist within and between:

- 1. Sectors: e.g., monogastric vs ruminant livestock (sections 2, 3, 4);
- 2. **Disciplines:** e.g., social, economic, environmental, biophysical, political, institutional etc. (section 5);
- 3. **Metrics:** e.g., GHG emissions per unit product, per unit protein, per unit animal etc. (sections 8, 9);
- 4. **Industry and value-chains**: e.g., primary producers, retailers, wholesalers, processors and consumers (section 8);
- 5. Scales: e.g., farm, region, country, continent, global (section 10), and
- 6. **Processes and systems**, considering holistic systems, rather than reductionist approaches that focus on one element while holding all others constant (section 8).

We are not advocating that future studies should be exhaustive, but rather that *future work give consideration to multi-dimensional aspects*. This could include computing alternative metrics of GHG emissions intensity; assessing the cost, environmental and social aspects of a given intervention; discussing the physical, financial and regional implications of adopting GHG emissions interventions; examining emissions leakage across industries, or quantifying trade-offs between the goals of farmers (often profitability rather than productivity) versus those of governments (to ensure productivity and thus food security).

Poore and Nemecek (2018) note that monitoring a single variable or proxy is often a poor predictor for net GHG emissions. They suggested that concurrent assessment of multiple impacts and avoiding proxies supported far better decisions, helping prevent harmful, unintended consequences. For example, adding N fertiliser to increase pasture biomass may result in pulses of N₂O that when cumulated can negate SOC benefit (Garnett et al., 2017). The environmental and social importance of different impacts varies locally, given land scarcity, endemic biodiversity and water quality, among other factors (Poore and Nemecek, 2018), suggesting multiple proxies are necessary to quantify key sustainability indicators in the social, environmental and economic dimensions.

We showed in Fig. 6 that assessments of social aspects of GHG emissions mitigation are particularly lacking. Salmon et al. (2018) suggested that consideration of the diverse objectives of stakeholders and agricultural paradigms should inform assessments of how the livestock sector in LMICs will evolve with future (and growing) demand for livestock products. Salmon et al. (2018) identified intensification trade-offs related to economic gains, gender equity, environmental concerns, human nutrition and food safety. They underscored the need to consider the distribution of benefits between *individuals* versus the average benefit is derived for a *population*.

Holistic, systems-based analyses are useful for examining feedback loops, leverage points, crossscale antagonisms or synergies and emergent properties (Alcock et al., 2015; Chang-Fung-Martel et al., 2017; Harrison et al., 2011a). Systems-based assessments can be used in a whole-farm sense (Ahmed et al., 2020; Phelan et al., 2015; Rawnsley et al., 2018) and across the value-chain to design interventions for reducing emissions associated with food loss and waste (Galford et al., 2020). Life cycle assessments are useful in analysing and attributing the emissions of a product from the 'cradle to grave' (or from resource extraction to waste disposal, see section 10). Because systems assessments enable analysis of interconnected processes, they are useful for determining leverage points at which a small change within a system could lead to larger changes in behaviours. Leverage points are thus one of the most critical points of the system for intervention (Weinberger et al., 2015).

System-based assessments can help bridge knowledge gaps across scales, from farm (Christie et al., 2018b; Harrison et al., 2014a; Harrison et al., 2014b; Pembleton et al., 2016) to regional and

continental levels (Chang et al., 2021). At the farm scale, soil carbon is often proposed as an avenue for GHG emissions mitigation (section 6). However, holistic assessment of other concurrent changes within a farm system may find that interventions to improve soil carbon can also increase whole farm emissions due to unexpected feedbacks wherein improvements in soil carbon raise pasture productivity (e.g. Rawnsley et al., 2018), leading to greater stocking rates and thus increase net GHG emissions at the farm scale (Fig. 8). Nevertheless, increased production intensity can provide opportunities for land sparing at farm or regional levels through displacement of less efficient production elsewhere, in which case they would be consistent with global climate change mitigation goals.

Holistic, systems-based assessments provide a pathway to *transcend multiple scales*. Chang et al. (2021) estimated regional patterns and global trends using a spatially-explicit land surface model and showed that human activities (rather than indirect effects of climate change) were responsible for causing managed grasslands to switch from a source to a sink of GHG emissions. The transition was mainly due to increased livestock numbers and accelerated conversion of natural lands to pasture. However, climate change also contributed to increased SOC sequestration as a result of increased productivity of grasslands due to increased atmospheric CO₂ and nitrogen deposition. Chang et al. (2021) showed that the net radiative forcing of global grasslands has been increasing since the 1960s but is currently close to neutral, because net global climate warming caused by managed grassland and ruminant livestock cancels the net climate cooling from carbon sinks. Systems-based approaches such as that used by Chang et al. (2021) provide an avenue for assessing the combined potential of multiple mitigation options imposed simultaneously (Box 2), facilitating insights into expected and unforeseen co-benefits and trade-offs.

9. Are sustainability metrics for livestock products appropriate, comparable and suitably cross-disciplinary?

Future research must shift from the prevailing and parochial view on carbon ('carbon myopia') towards more harmonised and globally-reconciled environmental, economic and social assessments. The derivation of sustainability methods and metrics in this way requires iterative input from multiple stakeholders, including farmers, industry and the scientific community, to civil society and governments at all levels. The application of holistic sustainability assessments would minimise risk that positive progress in one area (such as emissions mitigation) results in maladaptation in other dimensions, regions or sectors (such has diminished food security or natural habitat destruction).

As a first step towards more holistic sustainability comparisons, future work should quantify multiple metrics simultaneously: both within disciplines (alternative emissions metrics) and across disciplines, combining biophysical, social and economic indicators to examine livestock production from a social-ecological system perspective. The evolving science of global warming potential metrics is a case in point (Box 3): choice of metric for assessing climate impacts of CH₄ affects the level of responsibility for emissions reduction between countries, sectors and communities. This decision requires consideration of fairness, incentives and economic feedbacks, in addition to direct climate effects.

While there has been much work on the issue of GHG emissions allocation between products from the same animal (e.g. the partitioning of emissions between wool and meat for sheep, or between meat and milk for dairy cattle (e.g. Eady et al., 2012; Rice et al., 2017; Wiedemann et al., 2015)), there is less work on the appropriateness of emissions metrics for comparison of the relative GHG emissions intensities of livestock systems. This includes both the metric *per se* and the number of

metrics used to make any one comparison. By way of example, Cottle et al. (2016) modelled sheep GHG emissions intensities in 28 locations across Australia and showed that per animal (dry sheep equivalent), GHG emissions intensities were lower in cool temperate regions near the coast. In contrast, when emissions were computed per hectare, inland arid and semi-arid regions had lower emissions intensities. Such contradictions highlight a need to examine multiple metrics, including both emissions intensities and net emissions on a per land area and per animal basis. Emissions intensities are important when comparing production systems or enterprises (as in Cottle et al., 2016), and net emissions are important when calculating the relative effect of a production system on the atmosphere (and thus climate change mitigation potential).

Sustainability assessments across disciplines are also key to the development of more appropriate comparisons of the sustainability of agricultural systems going forwards. Evaluation of alternative emissions reduction strategies for livestock systems, and comparison with alternative livelihood systems, land uses and food production systems, should take an integrated systems approach (Bierbaum et al., 2018; O'Connell et al., 2019) that considers not just emissions but also environmental indicators, such as effects on soil acidification, water eutrophication, fresh water

Box 2: Combining or 'stacking' multiple GHG mitigation interventions

Although GHG mitigation options have received much attention in isolation, less work has been conducted on intervention 'stacking' (i.e., practices in which two or more GHG mitigation options are applied simultaneously). When applied holistically, concurrent assessment of multiple metrics can be used to evaluate the overall promise of 'bundled' or 'stacked' GHG mitigation options. For example, Harrison et al. (2016b) examined the nexus between GHG emissions, production and profitability for extensive beef farming systems in central Australia. They found that stacking of GHG mitigation options by mating maiden heifers earlier, optimising herd structure and increasing the fecundity of breeding cows resulted in higher gross margins than any of the three interventions applied individually. Adding the perennial legume leucaena (Leucaena leucephala) reduced gross margins but resulted in the lowest emissions intensity due to the positive effects of leucaena on liveweight gain and emissions mitigation (lower enteric methane and greater soil organic carbon sequestration). Similarly, Beukes et al. (2010) suggest that profitability of New Zealand dairy farms would be improved through the implementation of crossbred cows with higher genetic merit and reproductive performance, higher pasture quality, introduction of supplementary feeds (maize silage), and the application of a nitrification inhibitor under wet conditions to reduce N₂O emissions. Beauchemin et al. (2020) similarly indicate that a combination of mitigation strategies would be necessary to achieve substantial mitigation of enteric methane.

While the studies above examined the profitability of combined interventions, the costs of transitioning to a new system (e.g. to cows with greater genetic merit and/or reproductive performance) were not considered. Instead, the studies above examined the final steady-state scenarios after interventions were imposed. Given that significant and multi-faceted farming systems transitions can take time to be fully realised, such economic and biophysical considerations should be accounted for in future studies of this type.

Simultaneous improvements in production system (or business model) can be thought of as another example of intervention stacking. Steinfeld and Gerber (2010) concluded that in many LMIC, changing production systems from ruminants to monogastrics and continued efficiency gains in the production of feed and livestock could significantly attenuate the environmental impacts of livestock production systems. They indicated that although addressing excessive levels of consumption will help reduce environmental impact, there is a vast mitigation potential on the production side. Addressing environmental impacts of livestock on the production side may also carry important benefits for socially and economically disadvantaged livestock producers in LMIC.

Herrero et al. (2021) indicate that 'sociotechnical innovation bundles' (i.e. appropriately contextualised combinations of science and technology advancements coupled with institutional or policy adaptations that show promise for GHG mitigation) combined with institutional or policy reforms guided by a consistent mission or intentionality (Klerkx and Begemann, 2020) may be able to address multidisciplinary challenges and mitigate unintended consequences. Only with such concurrent policy reforms and other appropriate changes to the system will potential GHG mitigation be realised.

withdrawals and other ecosystem functions, as appropriate to context. In a globally reconciled and methodologically harmonised study, Poore and Nemecek (2018) showed that while GHG emissions intensities of lamb and mutton were higher than those of farmed crustaceans (10-25 kg cf. 5-15 CO₂-

e/100 g protein), effects of lamb/mutton production on acidification and eutrophication were much smaller than that of farmed crustaceans (50-75 *cf*. 50-150 g SO₂-e and 10-55 *cf*. 50-150 g $PO_4^{3-}e$ respectively). Similar parallels could be drawn with scarcity-weighted water withdrawals for beef, which in many cases were much lower than those of nuts and grains.

These examples suggest that if acidification and/or eutrophication were attributed a higher priority than GHG emissions in future assessments of sustainability, proponents for food production systems, policy and/or potentially societal preferences may shift favour towards red meat and away from farmed crustaceans. These policy shifts may then be at the expense of air quality (Domingo et al., 2021) and global warming if the net emissions from livestock were greater than those associated with crustaceans. Comparisons with other alternative protein sources, such as plant-based synthetic meat, microbial protein, and cultured meat, have similarly identified trade-offs between ER and inputs of energy, land, water and nutrients (Hu et al., 2020; Lynch and Pierrehumbert, 2019; Smetana et al., 2015; Spiller et al., 2020). *Consequential life cycle assessments* (CLCA) potentially represent a more integrated and arguably superior protocol for comparing sustainability dimensions of alternative protein sources as opposed to approaches that focus on carbon emissions alone.

'Wicked problems' of this type elicit questions on the priority placed on sustainability indicators, as a focus on one metric at one point in time (e.g., 'carbon myopia') could be at the detriment of another sustainability dimension in future (e.g., water quality, nutrient leaching, soil erosion, loss of biodiversity etc.). Similarly, changing priorities over time may shift societal and political preferences from one area of perceived importance to another, potentially negating years of previous effort and achievement in other sustainability dimensions. The priority placed on any sustainability indicator will be largely influenced by stakeholder interests. It is critically important that a representative range of stakeholders engage in constructive, iterative dialogue that results in the identification of new sustainability priorities, because perceived 'sustainability' will very much depend on the interests, background, education and potential benefit of those who contribute to formation of any new policy. This type of risk could be envisioned with a change of government that shifts their prevailing focus from carbon emissions mitigation to sustainable intensification; while enhanced food production and security may ensue, other societal outcomes may be maladaptive (for example, causing environmental degradation, financial instability, insecurity of land tenure or adverse human health outcomes) if they are not given appropriate and concurrent consideration.

We propose that future evaluation of emissions mitigation strategies for livestock systems includes comparison with alternative livelihoods, land uses and food production systems, applying an integrated systems approach (Bierbaum et al., 2018; O'Connell et al., 2019). Design of sustainable development interventions and governance systems should simultaneously consider *multiple objectives and metrics* to minimise trade-offs, stimulate co-benefits, enhance efficiency in policy implementation and promulgate more sustainable outcomes (Cowie et al., 2007; Tengberg and Valencia, 2018). In line with our results in Fig. 6, these sentiments suggest that greater understanding of social and economic contexts is critical to better inform landholder engagement and extension programs, and more effectively design policy incentives to enhance adoption of mitigation strategies.

Box 3: The state of the art in global warming metrics: are we comparing apples with apples?

Predominant GHGs in livestock systems include CO₂, CH₄, and N₂O. Methane is a powerful GHG but has a short atmospheric lifetime (~12 years) compared with 120 years for N₂O, and well over 500 years for CO₂ (Myhre et al., 2013). Following convention applied in national inventory reporting to the UNFCCC, global warming potential (GWP) has become an accepted metric for comparison of GHG emissions across agricultural products. GWP is calculated as the radiative forcing of a given GHG integrated over a chosen period (usually 100 years) expressed relative to the effect of an emission of CO₂ with units of CO₂-equivalents (CO₂-e).

While cumulative CO_2 emissions dominate anthropogenic warming over centuries, temperatures over the coming decades are also strongly influenced by short-lived climate pollutants (SLCP) including methane (Allen et al. 2018). Alternative GHG emission metrics have thus been proposed, including global temperature change (GTP) potential (Allen et al., 2016; Shine, 2009; Shine et al., 2007) and GWP* (Allen et al., 2018), which place alternative weighting on short-lived GHGs compared with GWP. GTP compares GHGs based on their effect on global mean surface temperate at the end of the chosen time horizon. GWP* compares CO_2 emissions to date with the current rate of emissions of SLCP, so to be computed requires two emissions measurements over a period of time (Δ t) – usually 20 years in the case of CH₄. Proponents argue that GWP* better reflects the shorter atmospheric lifetime but relatively higher radiative forcing of gases like CH₄, accounting the effects of both long-lived and short-lived pollutants on radiative forcing and temperatures over a wider range of time scales (Allen et al. 2018).

But GWP* also has limitations. The metric may penalise growth in livestock numbers, such as that anticipated in LMIC where livestock are critical for food security, and would give a negative CO₂-equivalent value in contexts where there is a static or declining livestock populations (e.g., Australia); the latter implying cooling, which is clearly not the case. The relevant application of GWP* to inform mitigation policy is thus contested, with concerns raised over equity implications (Harrison et al., 2014b; Rogelj and Schleussner, 2019).

Debate surrounding GHG equivalence reinforces the need for further development of metrics that facilitate more equitable approaches for comparing agricultural products – in particular, comparing short- and long-lived GHGs - but also in the context of offsets, for quantifying equivalence of carbon sequestration against GHGs with various lifetimes and differing impacts on the atmosphere. This is critically important, given the implications using GWP* may have in considering 'carbon neutral' or 'net zero emissions' targets for sectors emitting different GHG compositions (Lynch et al., 2020).

The development of such metrics would be expected to lead towards the derivation of functional units that could be applied to measure GHG emissions and account for pollution swapping, avoiding 'trivial solutions to a global problem' (Franks and Hadingham 2012).

10. In pursuit of multi-scale transdisciplinary solutions: consequential life cycle assessments and 'socio-economic planetary boundaries'

As discussed above, systems-based and holistic assessments may provide a prospective way forward for capturing multiple feedbacks within connected systems, including synergies, antagonisms and emergent properties. A systems-based approach frequently used to quantify carbon footprints associated with food production that accounts for some environmental trade-offs and co-benefits is *life-cycle assessment* (LCA; e.g. Alvarez-Hess et al., 2019; Beauchemin et al., 2010; Browne et al., 2011; Casey, 2005; Reijnders, 2012). Many 'conventional' or 'attributional' LCAs have shown that ruminant production has a GHG emissions intensity higher than that of crop production (Browne et al., 2011; Clune et al., 2017; Tuomisto and Teixeira de Mattos, 2011) and that livestock require several times the amount of land compared with production of vegetable proteins (Smith et al., 2013; Stehfest et al., 2009).

While product-focussed attributional LCAs show the relative impact of functionally-equivalent food products, they overlook the raison d'être of multifunctional land use systems, as well as the consequences of policy and consumer choice. For example, attributional LCA of dairy milk shows that subsistence dairy production in LMIC has higher emissions intensity than milk produced in a highly intensive dairy production system (e.g., compare the emissions intensity of African and European milk in Fig. 4). However, as explained above, livestock in smallholder systems serve many functions, including milk and meat production and draught power, but also provide less tangible social values, such as use for dowry, demonstration of prestige or wealth, and as key pillars of cultural identity (see Table 2 and Weiler et al., 2014). When GHG emissions were allocated between these various purposes, Weiler et al. (2014) found GHG emissions per unit weight of milk from smallholder dairying in Kenya of 2.0 (0.9-4.3) kg CO₂-e using economic allocation to food products only, 1.6 (0.8–2.9) kg CO₂-e when allocation was based on economic functions, and 1.1 (0.5–1.7) kg CO₂-e when emissions allocation considered the livelihood values of livestock. These emissions intensities are comparable to intensive dairy production systems (Alvarez-Hess et al., 2019; Beauchemin et al., 2010; Christie et al., 2018a; Galloway et al., 2018), clearly highlighting a need for careful consideration of the diverse economic, environmental and socio-economic roles that livestock play, particularly in smallholder systems.

Consequential life cycle assessment (CLCA) is emerging as a tool for capturing environmental impacts of production systems that go beyond physical relationships accounted for in conventional or attributional LCA (Earles and Halog, 2011). CLCA aims to describe how physical flows can change as a consequence of an increase or decrease in demand for the product under study. This is an important advancement on attributional LCA methods, particularly when applied to GHG emissions from the livestock sector in developing countries.

Consequential LCA has been used to assess emissions associated with consumer preferences in developed countries. Recent assessments have considered emissions associated with land use change resulting from dietary shifts in red meat consumption towards synthetic meat: there may be more pastoral and crop land available for reforestation in moving systems away from red meat, but conversely, an increase in crop production to supply the nutrients for plant-based meat substitutes could lead to long-term grasslands being cropped, losing substantial SOC and increasing N₂O emissions (Palmer et al., 2017; Pinto et al., 2004). A subsequent study showed that while cultured meat initially resulted in less climate warming than that from a cattle production system, the emissions gap narrowed in the long-term (Lynch and Pierrehumbert, 2019). The same study revealed that in some cases, cattle systems resulted in significantly less warming than cultured meat because

CH₄ emissions from ruminant production were short-lived, in contrast to the long-lived and fossilfuel derived CO₂. This insight again gives credence to the need for continuing investigation of alternative global warming metrics described in Box 3. A shift from livestock to cropping in LMIC may require more diesel use for mechanised cultivation as well as conventional fertilisers, increasing CO₂ and N₂O emissions. A transition away from mixed farming systems - particularly in subsistence agriculture – could also compromise local food security and resilience to climate change. While conventional LCA has been useful for exploring the comparative footprint of agricultural products, future policy would be better served by a more comprehensive and inclusive CLCA approach to better understand possible outcomes of policy choices (Brandão et al., 2017; Plevin et al., 2014).

A multi-scale approach broader than CLCA would be an expansion of the 'planetary boundaries' framework (Häyhä et al., 2016). The 'planetary boundaries' paradigm was developed assuming that transgression of one or more planetary boundaries could be deleterious and potentially catastrophic to humanity due to risks associated with crossing thresholds triggering non-linear, abrupt environmental tipping points within continental-scale to planetary-scale systems (Rockström et al., 2009). The framework defines a 'safe operating space for humanity', moving away from sectoral analyses of limits to growth aimed at minimising negative externalities, towards estimation of a safe space for human development. Planetary boundaries define, as it were, the boundaries of a "planetary playing field" for humanity if major global human-induced environmental change is to be avoided (Rockström et al., 2009). Two of the planetary boundaries have already been transgressed (biosphere integrity and biogeochemical flows; Fig. 9), with some authors suggesting that agriculture has been a major driver of this transgression (Campbell et al., 2017).

Planetary boundaries were originally conceived for nine key areas of the earth system (Fig. 9) but the work received criticism in failing to account for trade-offs between planetary boundaries (Lade et al., 2020). Expansion of the planetary boundaries to encompass socio-economic, cultural and ethical factors (Häyhä et al., 2016) would seem pertinent, as would approaches for accounting for interactions between planetary boundaries (Lade et al., 2020).

A greater focus on the development of food systems within a *'socio-economic planetary boundaries' systems-based paradigm* would be a fruitful endeavour for future comparisons of the sustainability of livestock production systems. Such focus may help shift the dialogue from a predominant and myopic focus on the technology aspects of GHG emissions mitigation to a multi-metric, multi-scale, transdisciplinary scenario analysis process. Development of such scenarios would help develop a positive visioning for the future, ensuring harmony across disciplines and global consistency in policy development.

The development of socio-economic planetary boundaries in this way should support and foster stakeholder dialogues, learning and understanding by visualising possible futures. Such scenarios would provide narratives to describe how a practice in a particular region may look (Weinberger et al., 2015) and could be used to develop a positive vision for future sustainable development.

11. The need for consistent, sustained and long-term research and development funding commitments for reducing livestock GHG emissions

Methane and nitrous oxide production are microbially-mediated processes that have evolved over 50M years. The development of mitigation strategies that fundamentally change these processes is thus a significant challenge, requiring sustained research commitment and funding continuity for

research groups addressing these issues. Investment for R&D to reduce agricultural emissions intensities has a high benefit: cost ratio, ultimately improving economic efficiencies, reducing both poverty and agricultural emissions (Laborde et al., 2021). Despite this, many countries operate with short-term, intermittent research and development (R&D) funding. In Australia for example, R&D investment in agricultural GHG emissions mitigation and climate change has historically been highly variable, resulting in recruitment then jettison of academic and institutional research capacity with each funding cycle. Agricultural R&D funding cycles in LMIC tends to be even more variable, often due to the short-term project-oriented nature of donor and development bank funding (Beintema et al., 2012). Such lack of continuing R&D investment has contributed to the paucity of GHG emissions mitigation technologies available for LMIC, as noted above. Considering the projected growth in livestock production and GHG emissions in LMIC in the next few decades, the absence of steady, sustained R&D funding heralds a dire outlook for enduring GHG emissions mitigation in developing nations.

12. Concluding remarks

The purpose of this review was to (1) examine the global distribution of livestock production and GHG emissions, (2) explore the co-benefits and trade-offs associated with economic, environmental and social implications of GHG emissions mitigation, (3) critique current approaches used to quantify and report livestock GHG emissions and (4), highlight opportunities for future GHG emissions mitigation research.

This review has helped distil many insights and opportunities for pathways forward:

- Globally, livestock production systems exist for several reasons, many of which are critical to livelihoods. In many areas, livestock are needed to satisfy a variety of human needs. In addition to production of meat, milk, eggs, wool, hides and skin, livestock provide draught power and nutrient cycling, supporting the environmental sustainability of production (Steinfeld et al., 2003). Ruminant livestock utilise non-arable land, converting fibrous and cellulosic materials into edible human protein. In many LMIC, livestock constitute the main (if not only) household capital reserve, serving as a strategic financial reserve that reduces risk and adds financial stability to the farming system (Steinfeld et al., 2003).
- 2. GHG emissions associated with ruminant livestock are greatest in LMIC. In contrast, the majority of GHG emissions mitigation research has been undertaken in and designed for developed countries. Currently, the developing world contributes 66% of global GHG emissions from ruminants: between 2017 and 2029, 80% of the growth in global meat production will occur in LMIC. The limited GHG emissions mitigation research and highly volatile R&D funding for such work in LMIC limits research capability and progress towards GHG emissions mitigation goals. Taken together, these trends underscore a dire need for research, development, extension and adoption of emissions mitigation technologies, skills and practices in LMIC. This is particularly so given the critical biophysical, cultural, financial and social dependencies of many smallholders on livestock.
- 3. The majority of GHG emissions mitigation research predominantly (in some cases myopically) focusses on technology aspects of GHG emissions mitigation. By corollary, there is a paucity of studies that concurrently assess social, economic and environmental aspects of GHG emissions mitigation. Of the 54 livestock GHG emissions mitigation interventions examined, only 16 had triple-bottom line benefits and medium-high mitigation potential. These included manure management, higher fecundity/weaning rates, increasing animal growth rates and reducing time

to slaughter. Planting deep-rooted legumes offers multiple avenues for emissions mitigation through enteric CH₄ mitigation as well as soil CS. The international scientific community must work to understand the wider social, economic and environmental co-benefits and trade-offs caused by the implementation of such interventions, because any change to existing practices could result in maladaptive outcomes, such as compromised food security or financial stability.

- 4. While multi-pronged efforts aimed at both emissions reduction (ER) and carbon sequestration (CS) are likely to achieve greater emissions mitigation, CS opportunities may provide greater potential for social, economic and environmental co-benefits. Strategic, thoughtful planting of trees and shelter-belts in areas of low pasture productivity paves avenues for CS but also heralds significant co-benefits, such as the provision of shade and shelter for livestock, habitat for native flora and fauna, fuel and firewood in LMIC and building of natural capital.
- 5. In addition to large areas of agroforestry that have already been adopted in LMIC, there appears to be considerable potential for further CS in soils and vegetation in LMIC. Despite the array of co-benefits that come with such emissions mitigation strategies, implementation of soil or vegetation measures specifically for GHG emissions mitigation through participation in carbon markets in LMIC has historically been hindered due to relatively high transaction costs and social barriers such as insecurity of land tenure. Improved adoption of practices for improving carbon stocks in soils and vegetation will require the implementation of both appropriate policy legislation and accompanying socio-economic developments (Kongsager et al., 2013).
- 6. The scientific community needs to shift from the frequent reductionist, unicentric focus on carbon, to more holistic, systems-based, multi-metric and transdisciplinary approaches. Such shifts will help uncover feedback loops, leverage points, cross-scale antagonisms/synergies and unforeseen emergent properties. Mainstream attention needs to progress from the CO₂-e mitigation lens in isolation to the investigation of multiple sustainability indicators that lead to profitable, practical and socially-acceptable solutions with cross-sectoral and multi-scale advantages. The derivation of harmonised, globally-reconciled sustainability metrics will require iterative dialogue from stakeholders at all levels, from farmers, industry and the scientific community to civil society and governments. The development of sustainability metrics in this way would be expected to limit cases in which positive progress in one area results in transgressive outcomes in other areas.
- 7. 'Wicked problems' in food sustainability elicit questions on how priority is placed on sustainability indicators, as a focus on one metric (e.g., carbon or GHG emissions) could be to the detriment of other sustainability metrics (e.g., water quality, nutrient leaching, soil erosion, loss of biodiversity etc.). Evolving priorities over time may shift societal and political preferences from one area of perceived importance to another. Shifts in industry priorities, governance and policy in this way could negate years of previous work in other areas. Risks of this type could be envisioned with a change to governments that are less interested in emissions mitigation and more focussed on sustainable intensification.
- 8. Nascent approaches such as consequential life cycle assessments (LCA) and socio-economic planetary boundaries provide opportunities to advance the state of the art of GHG emissions mitigation. In contrast to traditional attributional LCA, consequential LCA captures the wider environmental and socio-economic impacts of production systems, such as nutritional value, draught power, social status and financial capital provided by livestock systems in smallholder systems.
- 9. Integration of a system-based paradigm within a 'socio-economic planetary boundaries' framework provides another prospective pathway forward. This approach may yield fruitful comparisons of the sustainability of various livestock production systems and may help shift the dialogue from an individual technology and carbon emissions mitigation lens in isolation (our

phrase *carbon myopia*) towards a more nuanced global discussion, facilitating decision making through refinement of alternative scenarios, facilitating harmony across disciplines and consistency in global objectives.

13. Figure Legends

Fig. 1. Global GHG emissions from ruminant livestock (cattle, buffalo, sheep, and goat). Estimates were computed using IPCC Tier 1 Guidelines (IPCC, 2006) associated with on-farm emissions. Note different scaling on each ordinate axis. Values shown have been adapted from FAOSTAT (<u>http://www.fao.org/faostat</u>).

Fig. 2. Average meat supply per capita (kilograms per annum) versus gross domestic product (\$US GDP) per capita in 2015. Meat supply was computed as the total amount of the commodity available for human consumption during the reference period. Bubble size is proportional to population per country (China = 1,407 million, United States = 321 million and Ireland= 4.7 million). Values shown do not include fish or seafood (adapted from FAOSTAT http://www.fao.org/faostat/en/#data/FBS).

Fig. 3. Projected growth in global livestock meat production (carcass weight) in developed countries and LMIC between 2017 and 2029 (adapted from OECD, 2021).

Fig. 4. Disaggregated historical and future greenhouse gas emissions (GHG) associated with meat and milk production from key ruminant species (cow, sheep, goat and buffalo) for the main livestock producing continents from 1975 to 2050. X-axis values: 1 = Africa, 2 = Americas, 3 = Asia, 4 = Europe, 5 = Oceania. Values shown to the right of the dashed red line indicate future projected GHG emissions. Note differing scales on ordinate axes. Values shown were estimated using IPCC Tier 1 associated with on farm emissions (adapted from FAOSTAT http://www.fao.org/faostat).

Fig. 5. Emission intensities of meat and milk from key ruminants (cow, sheep, goat and buffalo) for the main livestock producing continents from 1975 to 2050. x-axis values: 1 = Africa, 2 = Americas, 3 = Asia, 4 = Europe, 5 = Oceania. Values shown were estimated using IPCC Tier 1 associated with on farm emissions. Note differing scales on ordinate axes (adapted from FAOSTAT <u>http://www.fao.org/faostat</u>).

Fig. 6. A. Number of Web of Science Core Collection documents published between 1945 and 2021 that include keywords 'greenhouse gas emissions mitigation and livestock' and/or additional biophysical, economic, environmental and social keywords; **B** corresponding search terms for LMIC or developing countries conducted using the search terms 'AND (LMIC OR (low AND Middle AND income) OR "developing countr*")'. Bar colours in **B** represent corresponding search colours in **A**; note change in order of histogram bars.

Fig. 7. Relationship between theoretical maximum and actual GHG mitigation potential as influenced by social (barriers to adoption, animal welfare, social licence etc.), political (carbon price, red tape, emission mitigation legislation etc), environmental (nutrient leaching, soil erosion, influence on biodiversity, ecosystems services etc.), economic (cost of implementation and relative return on investment etc.), and biophysical factors (climate, soil, geography, location etc). Social, political and environmental issues historically tend to be investigated last in studies of GHG emissions mitigation, whereas GHG emissions mitigation and productivity potentials are often evaluated first. Text adapted from Smith (2012).

Fig. 8 Example of a trade-off resulting from an intervention aimed at increasing soil carbon (e.g., through participation in an emissions trading scheme) that could result in higher net GHG emissions compared with business as usual. In the absence of external influence or constraints, landholders are likely to adapt grazing management to utilise any additional pasture or grassland production. This example illustrates the importance of assessing GHG emissions mitigation options holistically to manage unforeseen trade-offs, such as pollution swapping.

Fig. 9 Current status of the nine planetary boundaries. Below the green zone defines the safe operating space, yellow represents the uncertainty zone (increasing risk), red is the high-risk zone. The planetary boundary lies at the inner-most heavy circle. The control variable for climate change is equivalent atmospheric CO₂ concentration. Processes for which global-level boundaries are not quantified are shaded grey

(atmospheric aerosol loading, novel entities, and the functional role of biosphere integrity). Reproduced with permission from Steffen et al. (2015).

14. Tables

Table 1. Economic, environmental and social co-benefits and trade-offs associated with emissions reduction (ER) and/or carbon sequestration (CS) interventions in developed countries and LMICs. Economic co-benefits and trade-offs (Econ) consider productivity, profitability and opportunity costs. Environmental co-benefits (Envirn) were attributed based on air, water and land pollution, land degradation and risk of toxicity. Social aspects (Social) were assessed considering technology availability, social license, capacity for adoption, animal welfare, and public perceptions. Yellow shaded rows indicate GHG interventions applicable to LMIC and developed countries, white rows indicate interventions primarily applicable in developed countries. Within the nine categories, interventions are ranked in ascending order of positive economic, environmental and social effects. Legend (base of table): High \geq 30% mitigation; Med = 10-30% mitigation; Low = \leq 10% mitigation. Pos: positive; Neg: negative; NC: no change; '?' = unclear.

	Intervention	Co-benefits and trade-offs					Poforonco		
		Type Mitig		Econ	Envirn	Social	Reference		
1	1. Animal management or genetics								
	Transient confinement feeding of grazing animals to preserve ground cover and increase liveweight gain	ER, CS	••	=	↑	↓	(Cottle et al., 2016; Modernel et al., 2013; Molossi et al., 2020)		
	Genetic selection (residual feed intake) for low CH4 production	ER	•	ᡗ	Ţ	↓	(Alcock et al., 2015; Beauchemin et al., 2020; Bezerra et al., 2013; Hristov et al., 2013; Leahy et al., 2019)		
	Genetic selection for larger adult body size	ER	•	⊥	=	?	(Cottle et al., 2016)		
	Genetic selection for greater fleece weight production per animal	ER	•	<	=	=	(Alcock et al., 2015; Cottle et al., 2016)		
	Reducing age of first mating	ER	•	←	н	=	(Alcock et al., 2015; Christie et al., 2016; Cullen et al., 2016; Harrison et al., 2014a)		
	Extended seasonal lactation duration in dairy cows	ER	••	<	П	=	(Browne et al., 2014)		
	Recombination bovine somatotropin to increase growth rates	ER	•	<	Ţ	⊥	(Capper et al., 2008; Capper and Cady, 2012; Hristov et al., 2013; MacLeod and Moran, 2017)		
	Optimising herd structure for improved profit	ER	•	<	Ξ	1	(Harrison et al., 2016b)		
	Milking dairy cows less frequently	ER	•	↑	?	1	(Christie et al., 2014)		
	Improved animal health	ER	٠	1	1	↑	(Herrero et al., 2013b; Herrero et al., 2015; Shields and Orme-Evans, 2015)		
	Reduced adult and juvenile mortality at birth	ER	٠	↑	↑	↑	(Herrero et al., 2015; Hristov et al., 2013; Shields and Orme-Evans, 2015)		
	Reduced age at slaughter and days on feed	ER	••	1	↑	↑	(Herrero et al., 2015; Hristov et al., 2013; Shields and Orme-Evans, 2015)		
	Higher fecundity/higher weaning rates	ER	••	1	↑	↑	(Cullen et al., 2016; Harrison et al., 2014a; Harrison et al., 2014b; Ho et al., 2014)		

	Increased productivity/ growth rates of young livestock	ER	••	↑	1	Ŷ	(Beauchemin et al., 2020; Harrison et al., 2011a, b; Harrison et al., 2012a, b; Hristov et al., 2013; Leahy et al., 2019; Reisinger and Clark, 2018; Shields and Orme-Evans, 2015; Taylor and Eckard, 2016; Taylor et al., 2016)
	2. Feed additives and feeding manage	gement					•
	Nitrate feeding in rangeland environments	ER	•	\checkmark	\checkmark	\checkmark	(Beauchemin et al., 2020; Cottle et al., 2016; Herrero et al., 2015)
	Grape marc	ER	٠	≯	⊻	≯	(Cortés et al., 2020; Davison et al., 2020)
	Nitrification inhibitors	ER	••	\$	↓	→	(Herrero et al., 2015; Hristov et al., 2013; Kösler et al., 2019; Leahy et al., 2019)
	Tannins	ER	•	¥	1	→	(Beauchemin et al., 2020; Herrero et al., 2015)
	Chemical inhibitors (3- nitrooxypropanol)	ER	•••	?	↓	→	(Beauchemin et al., 2020; Herrero et al., 2015; Kösler et al., 2019; Leahy et al., 2019)
	Concentrates, e.g., grains (highly digestible feeds)	ER	••	↑	\checkmark	\$	(Beauchemin et al., 2020; Herrero et al., 2015; Hristov et al., 2013)
	Dietary lipids	ER	••	→	↑	↔	(Beauchemin et al., 2020; Herrero et al., 2015; Hristov et al., 2013; Ludemann et al., 2016)
	Rumen microbiome and fermentation	ER	••	?	\$	→	(Beauchemin et al., 2020; Hristov et al., 2013; Kumari et al., 2020; Leahy et al., 2019)
	Methane vaccine	ER	•••	?	=	\rightarrow	(Rolfe, 2001)
	Algal-derived	ER	•••	\$	₹	↑	(Beauchemin et al., 2020; Davison et al., 2020; Kinley et al., 2020; Roque et al., 2019)
	Feedstuffs with low N concentration	ER	٠	₹	⊥	₹	(Christie et al., 2014; Leahy et al., 2020; Reisinger and Clark, 2018)
	Biochar as animal feed supplement	ER	•	=	1	↑	(Roberts et al., 2010; Schmidt et al., 2019)
	3. Pasture types and management					-	T
	Birdsfoot trefoil (Lotus spp.)	ER, CS	•	→	1	=	(Doran-Browne et al., 2015)
	Biserrula	ER, CS	•••	\checkmark	1	\mathbf{r}	(Davison et al., 2020)
	Simplified/fewer intensive systems/lower fertiliser use	ER	•	\$	\$	\$	(Harrison et al., 2017a; Leahy et al., 2019; Reisinger and Clark, 2018)
	Silages	ER, CS	•••	\bigcirc	\bigcirc	\bigcirc	(Kumari et al., 2020)
	Pasture production improvement	ER	••	↑	↑	↑	(Alcock and Hegarty, 2006; Harrison et al., 2014a; Herrero et al., 2016; Hristov et al., 2013; Kumari et al., 2020; Smith et al., 2007)
	Fodder Beet	ER, CS	••	1	1	1	(Leahy et al., 2019; Ruis and Blanco- Canqui, 2017; Sun et al., 2016)
	Plantain	ER, CS	••	↑	1	1	(de Klein et al., 2020; Leahy et al., 2019; Luo et al., 2018)
	Leucaena	ER, CS	••	1	1	↑	(Harrison et al., 2015; Taylor and Eckard, 2016; Taylor et al., 2016; Tomkins et al., 2019)
L	Desmanthus	ER, CS	••	1	1	1	(Davison et al., 2020)
	Fodder Rape	ER, CS	•••	\uparrow	1	1	(Leahy et al., 2019; Ruis and Blanco-

						Canqui, 20	17; Sun et	al., 2016)	
4. Soil management and health									
Addition of phosphorus fertilisers	ER, CS	••	↑	¥	1	(Chan et al., 2010; Cottle et 2016; Harrison et al., 2014; al., 2015)		ttle et al., 2014a; Le	
Biochar to improve soil C	CS		\checkmark	1	=	(Joseph et	al., 2015)		
Converting annual crops to	CS	••	1	↑	↓	(Meier et a	al., 2020; N	leyer et al	
Improving soil carbon under trees	CS		=	^	^	(Doran-Bro	wne et al	2016)	
5 Manure management	0.5							, 2010)	
Solid liquid separation	ER	••	1	1	1	(Grossi et a 2013)	al., 2019; H	ristov et a	
Anaerobic digestion	ER	•••	1	1	1	, (Gerber an al., 2019; F	id Span, 20 Hristov et a	08; Grossi I., 2013)	
Decreased storage time	ER	•••	1	1	1	(Grossi et a 2013)	al., 2019; H	ristov et a	
requent manure removal	ER	•••	↑	↑	1	(Grossi et al., 2019; Hristov et 2013; Thornton, 2010)			
5. Agroforestry			T	T	1				
Planting trees on farm	CS	•••	↓	↑	\$	(Doran-Browne et al., 2016; al., 2019; Reisinger and Clark			
Forest conservation/Avoided deforestation	CS	•	Ţ	T	Ţ	(Petersen (Clark, 2018 2016)	et al., 2013 3; Rivera-Fe	; Reisinge erre et al.,	
7. Whole farm and regional manage	ement				•				
Integrated farming systems	ER, CS	••	1	1	?	(Thornton et al., 2018)			
Land restoration/Avoided land degradation	ER	••	↑	↑	1	(Doran-Browne et al., 2016; Herric et al., 2019; Rivera-Ferre et al., 2016)			
8. Technology and information serv	ices				1	,			
Renewable and alternate energy sources	CS	••	Ţ	1	=	(Fleming et al., 2019; Mayer et 2020)			
More accurate long-range and seasonal climate forecasts	ER	•	↑	=	↑	(Chang-Fung-Martel et al., 2017; Harrison et al., 2017b; Thornton et al., 2018)			
Digital services and decision support*	ER	••	1	1	1	(Fleming et al., 2019; Lovare 2020; Tullo et al., 2019)			
9. Social aspects									
Traditional/Indigenous knowledge	ER, CS	?	=	1	1	(Fleming et al., 2019; Kemp et al., 2013; Thornton et al., 2018)			
Cooperativism (Farmer producer organizations)	ER, CS	?	1	1	1	(Thornton	et al., 2018	3)	
*satellite imagery, big data, GPS, aut	tomation	, assess	ment ar	nd decisio	on suppor	rt.			
Legend: Low Med High	Ne	g N	eg/NC	NC	NC/Pos	Pos	Pos/Neg	Unc	
				_	^	•	^		
							-	· · · ·	

Table 2 Examples of economic, environmental and social co-benefits (black) and trade-offs (red) associated with farm GHG emissions mitigation in the livestock sector in developed countries and LMIC. Interventions that apply in both developed countries and LMIC are shown in parenthesis in the first column; all other interventions apply to developed countries.

Intervention examples		Economic	Environmental	Social
Feed additives	 3-NOP Algal derived Concentrates (some LMIC) Dietary lipids Grape marc Tannins (LMIC) Nitrate (rangelands) Low N feeds (LMIC) Biochar as a feed supplement 	 Some additives can improve liveweight gain and productivity Supplements that increase liveweight gain often lead to greater profitability Feeding biochar may be more cost effective than spreading on pastures, as livestock- spread biochar in faeces (Joseph et al., 2015) Nitrate as a feed additive can reduce productivity and therefore profitability Supplements that have no effect on productivity reduce profitability 	 Production of algal feeds can improve water quality depending on resource input and management, processing of spent water (NRC, 2013) Large scale production of algae may lead N, P and heavy metal pollution of environmental waters, leading to eutrophication The production and feeding of many additives may increase upstream (pre-farm) emissions, causing emissions leakage 	 Feeding biochar as livestock supplement often has positive effects on animal wellbeing through effects on toxin absorption, digestion, blood parameters, feed efficiency and meat quality (Schmidt et al., 2019) Feeding excessive tannins impacts on animal performance and induces metabolic disorders (animal welfare) (Jerónimo et al., 2016) Feeding concentrates to animals increases competition for arable land for food production Some feed additives may adversely impact animal health (e.g. possible carcinogenic effects of algal-derived feeds) or cause undesirable residues in animal products (tannins and saponins)(Martin et al., 2010)
Animal genetics	 Selection for low CH4 (low residual feed intake) Selection for larger body size (LMIC) Greater fleece weight Higher fecundity breeds (LMIC) Rumen microbiome and CH4 fermentation manipulation Livestock cloning (Godfray et al., 2010) 	 High fecundity breeds often lead to greater liveweight production More profitable (fertile) bulls can produce daughters that yield less milk but this can be more than compensated for by reduced health costs and lower herd replacement rates Breeding costs can be high due to the long lead times of such programmes Larger and faster-growing animals require more feed, which is not always available in smallholder systems – may be negative economic impacts where farmers are unable to provide adequate inputs to support reproduction and liveweight gain 	 Higher feed conversion efficiency animals require less feed per unit liveweight gain and less feed overall, reducing pressure on pastures Breeding may lead to higher ground cover, reduced erosion and greater enteric CH₄ production if stocking rates are lowered 	 Higher fecundity breeds of sheep often have higher lamb mortalities at birth because twins/triplets have lower body weight and are weaker (animal welfare implications) Breeding for milk yield or liveweight gain in isolation can be correlated with reduced fertility and health traits (Lawrence et al., 2004)
An im al	Extended lactation (LMIC)Reduced age of first mating	Reducing milking frequency can improve profitability through reduced labour costs	 Reduced milking frequency can cause less pasture trampling (pugging) damage 	 Milking cows less frequently can reduce labour inputs, reduce worker

	 (LMIC) Optimising herd structure Recomb. bovine somatotropin Milking cows less frequently Improved animal health (LMIC) Reduced mortality (LMIC) Improved weaning rates (LMIC) Seasonal confinement feeding Increased productivity of young livestock (LMIC) 	 but with similar milk income Optimising herd structure and incr growth rates of young animals can significantly raise profitability Improving weaning rates/fecundit animal husbandry often leads to g profitability Practices that rely on improving pr through improvements in liveweig may be at a disadvantage if marke decrease 	reasing h ty through greater rofit ght gain et prices	Confinement feeding can improve ground cover and reduce soil erosion, protect soil C Increasing stocking rates in LMIC increases competition between domesticated and wild livestock for scarce resources (Thornton and Gerber, 2010) Optimising herd structure can increase stocking rates, enteric CH ₄ and soil erosion (Thornton and Gerber, 2010)	 fatigue and improve wellbeing Reducing milking frequency can improve animal welfare though improved hoof health and fewer hoof diseases Earlier mating results in lower conception rates at first mating
Pasture types	 Silages (LMIC) Fodder Beet Plantain Leucaena (LMIC) Desmanthus (LMIC) Fodder Rape Birdsfoot trefoil (LMIC) Biserrula (LMIC) 	 Establishing new pastures can lead greater digestibility per area and seasonally, leading to greater livev gains and profitability Establishing new pasture types ma extensive capital investment (i.e., outlay) 	d to weight ay require capital	 Increased use of legumes reduces needs for synthetic N, lowering N2O emissions Deep-rooted pastures sequester additional soil C at depth, increasing soil organic CS per unit area Legumes contain anti-methanogenic compounds, preventing enteric CH4 Perennial crops used for energy production can favour biodiversity if they displace annual crops (Berndes and Börjesson, 2002) More productive forages allow higher stocking rate which in turn facilitates higher enteric CH4 per unit area (Fig. 8) Legumes can cause soil acidification Some introduced pastures can become noxious weeds (Lonsdale, 1994) 	 Using legumes provides a source of higher dietary crude protein, increasing livestock performance and welfare
Pasture management	 Addition of N fertilisers (some LMIC) Addition of P fertilisers (some LMIC) Converting annual crops to permanent pastures (LMIC) Nitrification inhibitors Simplified farming systems/lower fertiliser use 	 Access to some ecosystems service markets (e.g. improved water qual increase Where fertility is sub-optimal and the marginal return is greater than marginal cost, fertiliser addition ca improved production and profit ou Reducing input costs through redu 	where n the an lead to utcomes	Biochar addition can increase soil C Deep-rooted legumes can sequester C at greater depths, fostering greater sequestration per area Converting to permanent pastures (from annual crops) can improve soil carbon, increase water infiltration, reduce soil erosion and improve drought tolerance (Langworthy et al., 2018)	 Practices leading to lower fertiliser and pesticide use may improve health of surrounding communities and nitrates in drinking water Conversion of cropping land to permanent pastures may cause GHG emissions leakage, in which other land is used for crop production

	•	(LMIC) Addition of biochar	•	fertiliser inputs may reduce profitability In some cases, nitrification inhibitors are prohibitively expensive	•	N fertiliser can cause soil acidification, pollution of waterways and eutrophication Reducing fertiliser use reduces fertility and can result in loss of soil carbon	•	High reliance on synthetic fertilisers of some systems can be seen by the public as unsustainable, e.g., excessive use of rock phosphate fertilisers
					•	Some pastures may require higher pesticide and herbicide usage		
Manure management	•	Solid liquid separation Anaerobic digestion (LMIC) Decreased storage time Frequent manure removal (LMIC) Conversion to biochar	•	Recycled manures can be used as fuel sources Spreading of manure on pastures can improve profitability	• • • •	Increased recovery of nutrients, organic materials Lower nutrient leaching into waterways Improved nutrient-use efficiencies Reduced need for synthetic fertilisers (Cohen et al., 2021) Reduced ammonia volatilisation (Smith et al., 2021) Addition of manures to pastures can pollute waterways	•	Manures spread onto land used for crop production may contaminate grain, having human health implications

orestry and vegetation management	•	Planting trees on farm, improved soil C under trees (LMIC) Forest conservation (LMIC) Avoided deforestation (LMIC)	•	Agroforestry may increase access to some ecosystems services markets (water and air quality, biodiversity, reforestation, afforestation) (Bustamante et al., 2014) Income diversification and improved profit through timber production reduces reliance on animal products for income Afforestation increases access to firewood in LMIC Some tree species used for agroforestry provide high quality supplements for cattle Reduced land clearing can increase financial hardship in some communities Planting trees on productive lands may reduce profit due to loss of pasture	•	Increased biodiversity in soils and in trees Improved soil organic carbon Conservation of native forests (Cohen et al., 2021) Permanence of carbon (Cohen et al., 2021) Afforestation can increase biodiversity, both flora and fauna (e.g. wildlife corridors) Perennial vegetation strips provide shelter to livestock, reduce wind erosion and control dryland salinity Agroforestry may increase risk of fires Planting trees on productive land can result in loss of pasture area and lower livestock production. This could lead to conversion of forest to pastures, causing leakage Habitat provision for noxious weeds, pests and vermin (e.g rabbits) which may cause damage to not only natural areas but also	•	Improved consumer perceptions of environmental and land stewardship Forestry, afforestation and reforestation can provide the impetus to shift from fossil-fuel dependent societies to those more focussed on environmental and green energies Afforestation and increased firewood supply can reduce poverty in LMIC GHG emissions leakage if avoided deforestation causes other land to be used for crop or pasture production Forestry can compete with land available for grazing
rofo						improved pastures		
Ag					•	Planting trees in natural grasslands may contribute to losses of unique biodiversity		
						and changes to water movement through		
						the landscape (Abreu et al., 2017; Honda and Durigan, 2016)		
	•	Prevention/cessation of	•	Techniques that reduce or change the	•	Soil organic CS potential for improvement on	•	Renewable energy can contribute to
з		stubble or grass burning		seasonal timing of burning can significantly		degraded landscapes is often very high (Lal,		poverty alleviation in LMIC
ner		(LMIC)		reduce carbon lost to the atmosphere		2004; Pang et al., 2019)	•	Avoiding land degradation improves
gen	•	Integrated farming systems	•	Avoided land degradation increases arable	•	Reducing practices that use burning prevent		public perception of farmer
nag		(LMIC)		area and most often profitability		carbon loss to the atmosphere, conserve		environmental stewardship
na	•	(I MIC)	•	Land acquisition and enterprise		erosion through retaining surface organic		Use of energy crops can compete with
l pr	•	Land restoration (LMIC)		efficiencies can lower costs		matter		land available for grazing, reducing
lar	•	Avoided land degradation			•	Integrated farming systems lower reliance on		food security
pu		(LMIC)	•	Expanding enterprise or farm size may		fossil fuels and grid-based power generation,		
ла	•	Renewable and alternate		require significant capital investment and		improve pest management and biodiversity		
arr		energy sources		long-term debt		(Conen et al., 2021)		
le f	•	Further land acquisition			•	Use of alternative fuels/feedstocks reduce		
ho	•	(Iarm Size expansion, LIVIIC)				renance on synthetic rentilisers		
3	•	feedstocks (I MIC)			•	Use of alternative fuels may cause pollution		
						swapping (e.g., switching from petroleum to		

	More accurate long range	Increased access to digital convisos	 ethanol may increase ozone pollution) Agricultural land expansion removes native habitats, reduces soil fertility and biodiversity Higher growth rates of young animals may lead to higher stocking rates and emissions (Fig. 8) 	Digital accors will enable increased
IT services	 More accurate forg-range and seasonal climate forecasts (LMIC) Digital services and decision support (LMIC) 	 Increased access to digital services increases timeliness of management and reduces labour costs (Ara et al., 2021) Some food technologies have higher energy demand, resulting in higher input costs 	 Greater use of technology facilitates remote management, reducing emissions associate with manual labour and travel for livestock management and transport New satellite imagery sensors can be used to detect and prevent illegal logging, preventing CO₂ emissions before they occur (Curry and Harris, 2019; Jones et al., 2020) 	 Digital access will enable increased access to and awareness of information in some regions Prevention of illegal logging with satellite imagery improves human health and wellbeing (Jones et al., 2020) Digital services/internet access may not be available in some rural areas, particularly in LMIC
Social and policy enablers	 Traditional/Indigenous knowledge Co-operativism (Farmer producer organizations, LMIC) Carbon aggregation across farms 	 'Carbon aggregation' (where multiple farms participate as one in emissions mitigation) can reduce compliance costs and increase profit of individual farms Livestock may be the only income source for some producers in LMIC through food and non-food products (meat, milk, wool, hides, skin) Livestock can serve as financial instruments, providing LMIC households with an option for accumulating capital 	 The majority of life-cycle analyses focus on one emissions metric and are not consequential LCA (section 10) Environmental stewardship practices enacted may depend on social norms, access to information and available existing legislation that reduces emissions sustainably 	 GHG mitigations interventions often (but not always) entail net benefits for public health (Haines et al., 2009) In parts of Africa, social relationships and status are partly defined by the size of the household's livestock holding In some LMIC, livestock sharing strengthens social relationships through use as a dowry/bride price, and as allocations to family members as loans (Kitalyi et al., 2005) Livestock contribute to food security and provide sustenance to many undernourished people in LMIC, particularly children (Neumann et al., 2003) Policy incentives for C aggregation may not exist or can be difficult to co- ordinate

15. Acknowledgements

We thank Agri-Benchmark and the Thünen Institute of Farm Economics for allowing access to and publication of their global dataset for GHG emissions, animal numbers and productivity. This work was funded by the Tasmanian Institute of Agriculture, the University of Tasmania, the University of Melbourne, the Commonwealth Scientific Industrial Research Organisation (CSIRO) and the MLA Donor Company as part of the Nexus project for profitable, sustainable livestock businesses in an increasingly variable climate (P.PSH.1219, P.PSH.1236 and P.PSH.1248).

16. References

Abbott, D.W., Aasen, I.M., Beauchemin, K.A., Grondahl, F., Gruninger, R., Hayes, M., Huws, S., Kenny, D.A., Krizsan, S.J., Kirwan, S.F., Lind, V., Meyer, U., Ramin, M., Theodoridou, K., von Soosten, D., Walsh, P.J., Waters, S., Xing, X. (2020) Seaweed and Seaweed Bioactives for Mitigation of Enteric Methane: Challenges and Opportunities. Animals 10.

Abernethy, B., Rutherford, I. (1999) Guidelines for stabilising stream banks with riparian vegetation. Cooperative Research Centre for Catchment Hydrology. Department of Geography and Environmental Studies, University of Melbourne, Parkville, Victoria, pp. 1-137.

Abreu, R.C., Hoffmann, W.A., Vasconcelos, H.L., Pilon, N.A., Rossatto, D.R., Durigan, G.J.S.a. (2017) The biodiversity cost of carbon sequestration in tropical savanna. 3, e1701284.

Ahmed, M., Ahmad, S., Waldrip, H.M., Ramin, M., Raza, M.A. (2020) Whole Farm Modeling: A Systems Approach to Understanding and Managing Livestock for Greenhouse Gas Mitigation, Economic Viability and Environmental Quality. In Animal Manure (eds H. Waldrip, P. Pagliari and Z. He). <u>https://doi.org/10.2134/asaspecpub67.c25</u>.

Alcock, D., Hegarty, R.S. (2006) Effects of pasture improvement on productivity, gross margin and methane emissions of a grazing sheep enterprise. International Congress Series 1293, 103-106.

Alcock, D.J., Harrison, M.T., Rawnsley, R.P., Eckard, R.J. (2015) Can animal genetics and flock management be used to reduce greenhouse gas emissions but also maintain productivity of wool-producing enterprises? Agricultural Systems 132, 25-34.

Alexander, P., Brown, C., Arneth, A., Finnigan, J., Rounsevell, M.D.A. (2016) Human appropriation of land for food: The role of diet. Global Environmental Change 41, 88-98.

Alexander, P., Rounsevell, M.D.A., Dislich, C., Dodson, J.R., Engström, K., Moran, D. (2015) Drivers for global agricultural land use change: The nexus of diet, population, yield and bioenergy. Global Environmental Change 35, 138-147.

Allen, M.R., Fuglestvedt, J.S., Shine, K.P., Reisinger, A., Pierrehumbert, R.T., Forster, P.M. (2016) New use of global warming potentials to compare cumulative and short-lived climate pollutants. Nature Climate Change 6, 773-776.

Allen, M.R., Shine, K.P., Fuglestvedt, J.S., Millar, R.J., Cain, M., Frame, D.J., Macey, A.H. (2018) A solution to the misrepresentations of CO2-equivalent emissions of short-lived climate pollutants under ambitious mitigation. npj Climate and Atmospheric Science 1, 16.

Alvarez-Hess, P.S., Little, S.M., Moate, P.J., Jacobs, J.L., Beauchemin, K.A., Eckard, R.J. (2019) A partial life cycle assessment of the greenhouse gas mitigation potential of feeding 3-nitrooxypropanol and nitrate to cattle. Agricultural Systems 169, 14-23.

Andersson, K., Lawrence, D., Zavaleta, J., Guariguata, M.R. (2016) More Trees, More Poverty? The Socioeconomic Effects of Tree Plantations in Chile, 2001–2011. Environmental Management 57, 123-136.

Ara, I., Harrison, M.T., Whitehead, J., Waldner, F., Bridle, K., Gilfedder, L., Marques da Silva, J., Marques, F., Rawnsley, R. (2021) Modelling seasonal pasture growth and botanical composition at the paddock scale with satellite imagery. in silico Plants 3.

Badgery, W.B., Mwendwa, J.M., Anwar, M.R., Simmons, A.T., Broadfoot, K.M., Rohan, M., Singh, B.P. (2020) Unexpected increases in soil carbon eventually fell in low rainfall farming systems. Journal of Environmental Management 261, 110192.

Badgery, W.B., Simmons, A.T., Murphy, B.W., Rawson, A., Andersson, K.O., Lonergan, V.E. (2014) The influence of land use and management on soil carbon levels for crop-pasture systems in Central New South Wales, Australia. Agriculture, Ecosystems & Environment 196, 147-157.

Bai, Z., Ma, W., Ma, L., Velthof, G.L., Wei, Z., Havlík, P., Oenema, O., Lee, M.R., Zhang, F.J.S.a. (2018) China's livestock transition: Driving forces, impacts, and consequences. 4, eaar8534.

Bajželj, B., Richards, K.S., Allwood, J.M., Smith, P., Dennis, J.S., Curmi, E., Gilligan, C.A. (2014) Importance of food-demand management for climate mitigation. Nature Climate Change 4, 924-929.

Bardi, U., El Asmar, T., Lavacchi, A. (2013) Turning electricity into food: the role of renewable energy in the future of agriculture. Journal of Cleaner Production 53, 224-231.

Bayala, J., Prieto, I. (2020) Water acquisition, sharing and redistribution by roots: applications to agroforestry systems. Plant and Soil 453, 17-28.

Beauchemin, K.A., Henry Janzen, H., Little, S.M., McAllister, T.A., McGinn, S.M. (2010) Life cycle assessment of greenhouse gas emissions from beef production in western Canada: A case study. Agricultural Systems 103, 371-379.

Beauchemin, K.A., Ungerfeld, E.M., Eckard, R.J., Wang, M. (2020) Review: Fifty years of research on rumen methanogenesis: lessons learned and future challenges for mitigation. Animal 14, s2-s16.

Beier, C., Beierkuhnlein, C., Wohlgemuth, T., Penuelas, J., Emmett, B., Körner, C., de Boeck, H., Christensen, J.H., Leuzinger, S., Janssens, I.A., Hansen, K. (2012) Precipitation manipulation experiments – challenges and recommendations for the future. Ecology Letters 15, 899-911.

Beintema, N., Stads, G.-J., Fuglie, K., Heisey, P. (2012) ASTI global assessment of agricultural R&D spending - Developing countries accelerate investment. Agricultural Science and Technology Indicators, Rome Italy, 1-24.

Bell, L.W., Harrison, M.T., Kirkegaard, J.A. (2015) Dual-purpose cropping – capitalising on potential grain crop grazing to enhance mixed-farming profitability. Crop and Pasture Science 66, i-iv.

Bell, M.J., Eckard, R.J., Harrison, M.T., Neal, J.S., Cullen, B.R. (2013) Effect of warming on the productivity of perennial ryegrass and kikuyu pastures in south-eastern Australia. Crop and Pasture Science 64, 61-70.

Bellarby, J., Tirado, R., Leip, A., Weiss, F., Lesschen, J.P., Smith, P. (2013) Livestock greenhouse gas emissions and mitigation potential in Europe. Global Change Biology 19, 3-18.

Berndes, G., Börjesson, P. (2002) Multi-functional biomass production systems. Available: <u>http://www.elkraft.ntnu.no/eno/konf_pub/ISES2003/full_paper/6%20MISCELLANEOUS/06%204.pdf</u> [Accessed 31 May 2006].

Beukes, P.C., Gregorini, P., Romera, A.J., Levy, G., Waghorn, G.C. (2010) Improving production efficiency as a strategy to mitigate greenhouse gas emissions on pastoral dairy farms in New Zealand. Agriculture, Ecosystems & Environment 136, 358-365.

Bezerra, L.R., Sarmento, J.L.R., Neto, S.G., de Paula, N.R.O., Oliveira, R.L., do Rêgo, W.M.F. (2013) Residual feed intake: a nutritional tool for genetic improvement. Tropical Animal Health and Production 45, 1649-1661.

Bierbaum, R., Cowie, A., Barra, R., Ratner, B., Sims, R., Stocking, M., Durón, G., Leonard, S., Whaley, C.J.S., Technical Advisory Panel to the Global Environment Facility. Washington, D. (2018) Integration: To solve complex environmental problems. https://www.thegef.org/sites/default/files/publications/STAP%20Report%20on%20integration.PDF.

Bogie, N.A., Bayala, R., Diedhiou, I., Conklin, M.H., Fogel, M.L., Dick, R.P., Ghezzehei, T.A. (2018) Hydraulic Redistribution by Native Sahelian Shrubs: Bioirrigation to Resist In-Season Drought. 6.

Bossio, D.A., Cook-Patton, S.C., Ellis, P.W., Fargione, J., Sanderman, J., Smith, P., Wood, S., Zomer, R.J., von Unger, M., Emmer, I.M., Griscom, B.W. (2020) The role of soil carbon in natural climate solutions. Nature Sustainability 3, 391-398.

Bouwman, A.F., Van Vuuren, D.P., Derwent, R.G., Posch, M. (2002) A Global Analysis of Acidification and Eutrophication of Terrestrial Ecosystems. Water, Air, and Soil Pollution 141, 349-382.

Brandão, M., Martin, M., Cowie, A., Hamelin, L., Zamagni, A., (2017) Consequential life cycle assessment: what, how, and why?, Encyclopedia of Sustainable Technologies. Elsevier, pp. 277-284.

Brondz, I. (2018) Why Judaism and Islam Prohibit Eating Pork and Consuming Blood as a Food? Scientific Research 4.

Browne, N.A., Behrendt, R., Kingwell, R.S., Eckard, R.J. (2014) Does producing more product over a lifetime reduce greenhouse gas emissions and increase profitability in dairy and wool enterprises? Animal Production Science 55, 49-55.

Browne, N.A., Eckard, R.J., Behrendt, R., Kingwell, R.S. (2011) A comparative analysis of on-farm greenhouse gas emissions from agricultural enterprises in south eastern Australia. Animal Feed Science and Technology 166-167, 641-652.

Bustamante, M., Robledo-Abad, C., Harper, R., Mbow, C., Ravindranat, N.H., Sperling, F., Haberl, H., de Siqueira Pinto, A., Smith, P. (2014) Co-benefits, trade-offs, barriers and policies for greenhouse gas mitigation in the agriculture, forestry and other land use (AFOLU) sector. Global Change Biology 20, 3270-3290.

Byrnes, R.C., Eastburn, D.J., Tate, K.W., Roche, L.M. (2018) A Global Meta-Analysis of Grazing Impacts on Soil Health Indicators. Journal of Environmental Quality 47, 758-765.

Cacho, O.J., Marshall, G.R., Milne, M. (2005) Transaction and abatement costs of carbon-sink projects in developing countries. Environment and Development Economics 10, 597-614.

Callaghan, M.J., Tomkins, N.W., Benu, I., Parker, A.J. (2014) How feasible is it to replace urea with nitrates to mitigate greenhouse gas emissions from extensively managed beef cattle? Animal Production Science 54, 1300-1304.

Callaghan, M.J., Tomkins, N.W., Hepworth, G., Parker, A.J. (2021) The effect of molasses nitrate lick blocks on supplement intake, bodyweight, condition score, blood methaemoglobin concentration and herd scale methane emissions in *Bos indicus* cows grazing poor quality forage. Animal Production Science 61, 445-458.

Campbell, B.M., Beare, D.J., Bennett, E.M., Hall-Spencer, J.M., Ingram, J.S.I., Jaramillo, F., Ortiz, R., Ramankutty, N., Sayer, J.A., Shindell, D. (2017) Agriculture production as a major driver of the Earth system exceeding planetary boundaries. Ecology and Society 22.

Capper, J., Cady, R., Bauman, D., (2008) Increased production reduces the dairy industry's environmental impact, Preceedings of the Cornell Nutrition Conference fot Feed Manufactturerst. Syracuse, New York. Citeseer.

Capper, J.L., Cady, R.A. (2012) A comparison of the environmental impact of Jersey compared with Holstein milk for cheese production1. Journal of Dairy Science 95, 165-176.

Caro, D., Davis, S.J., Bastianoni, S., Caldeira, K. (2014) Global and regional trends in greenhouse gas emissions from livestock. Climatic Change 126, 203-216.

Caro, D., Davis, S.J., Kebreab, E., Mitloehner, F. (2018) Land-use change emissions from soybean feed embodied in Brazilian pork and poultry meat. Journal of Cleaner Production 172, 2646-2654.

Casey, J.W.H.N.M. (2005) The relationship between greenhouse gas emissions and the intensity of milk production in Ireland. Journal of Environmental Quality. 3, 429-436.

Cerdà, A., Rodrigo-Comino, J. (2020) Is the hillslope position relevant for runoff and soil loss activation under high rainfall conditions in vineyards? Ecohydrology & Hydrobiology 20, 59-72.

Chan, K.Y., Oates, A., Li, G.D., Conyers, M.K., Prangnell, R.J., Poile, G., Liu, D.L., Barchia, I.M. (2010) Soil carbon stocks under different pastures and pasture management in the higher rainfall areas of south-eastern Australia. Soil Research 48, 7-15.

Chang-Fung-Martel, J., Harrison, M.T., Rawnsley, R., Smith, A.P., Meinke, H. (2017) The impact of extreme climatic events on pasture-based dairy systems: a review. Crop and Pasture Science 68, 1158-1169.

Chang, J., Ciais, P., Gasser, T., Smith, P., Herrero, M., Havlík, P., Obersteiner, M., Guenet, B., Goll, D.S., Li, W., Naipal, V., Peng, S., Qiu, C., Tian, H., Viovy, N., Yue, C., Zhu, D. (2021) Climate warming from managed grasslands cancels the cooling effect of carbon sinks in sparsely grazed and natural grasslands. Nature Communications 12, 118.

Christie, K.M., Harrison, M.T., Trevaskis, L.M., Rawnsley, R.P., Eckard, R.J. (2016) Modelling enteric methane abatement from earlier mating of dairy heifers in subtropical Australia by improving diet quality. Animal Production Science 56, 565-573.

Christie, K.M., Rawnsley, R.P., Harrison, M.T., Eckard, R.J. (2014) Using a modelling approach to evaluate two options for improving animal nitrogen use efficiency and reducing nitrous oxide emissions on dairy farms in southern Australia. Animal Production Science 54, 1960-1970.

Christie, K.M., Rawnsley, R.P., Phelps, C., Eckard, R.J. (2018a) Revised greenhouse-gas emissions from Australian dairy farms following application of updated methodology. Animal Production Science 58, 937-942.

Christie, K.M., Smith, A.P., Rawnsley, R.P., Harrison, M.T., Eckard, R.J. (2018b) Simulated seasonal responses of grazed dairy pastures to nitrogen fertilizer in SE Australia: Pasture production. Agricultural Systems 166, 36-47.

Clune, S., Crossin, E., Verghese, K. (2017) Systematic review of greenhouse gas emissions for different fresh food categories. Journal of Cleaner Production 140, 766-783.

Cockfield, G., Shrestha, U., Waters, C. (2019) Evaluating the potential financial contributions of carbon farming to grazing enterprises in Western NSW. The Rangeland Journal 41, 211-223.

Cohen, B., Cowie, A., Babiker, M., Leip, A., Smith, P. (2021) Co-benefits and trade-offs of climate change mitigation actions and the Sustainable Development Goals. Sustainable Production and Consumption 26, 805-813.

Cohn, A.S., Mosnier, A., Havlík, P., Valin, H., Herrero, M., Schmid, E., O'Hare, M., Obersteiner, M. (2014) Cattle ranching intensification in Brazil can reduce global greenhouse gas emissions by sparing land from deforestation. Proceedings of the National Academy of Sciences 111, 7236.

Conant, R.T., Paustian, K. (2002) Potential soil carbon sequestration in overgrazed grassland ecosystems. Global Biogeochemical Cycles 16, 90-91-90-99.

Cortés, A., Moreira, M.T., Domínguez, J., Lores, M., Feijoo, G. (2020) Unraveling the environmental impacts of bioactive compounds and organic amendment from grape marc. Journal of Environmental Management 272, 111066.

Cottle, D.J., Harrison, M.T., Ghahramani, A. (2016) Sheep greenhouse gas emission intensities under different management practices, climate zones and enterprise types. Animal Production Science 56, 507-518.

Cowie, A., Schneider, U.A., Montanarella, L. (2007) Potential synergies between existing multilateral environmental agreements in the implementation of land use, land-use change and forestry activities. Environmental Science & Policy 10, 335-352.

Cowie, A.L., Waters, C.M., Garland, F., Orgill, S.E., Baumber, A., Cross, R., O'Connell, D., Metternicht, G. (2019) Assessing resilience to underpin implementation of Land Degradation Neutrality: A case study in the rangelands of western New South Wales, Australia. Environmental Science & Policy 100, 37-46.

Cullen, B.R., Eckard, R.J., Timms, M., Phelps, D.G. (2016) The effect of earlier mating and improving fertility on greenhouse gas emissions intensity of beef production in northern Australian herds. The Rangeland Journal 38, 283-290.

Curry, J., Harris, N. (2019) Powering the Environmental Internet of Things. Sensors 19.

Davison, T.M., Black, J.L., Moss, J.F. (2020) Red meat—an essential partner to reduce global greenhouse gas emissions. Animal Frontiers 10, 14-21.

de Klein, C.A., van der Weerden, T.J., Luo, J., Cameron, K.C., Di, H. (2020) A review of plant options for mitigating nitrous oxide emissions from pasture-based systems. New Zealand Journal of Agricultural Research 63, 29-43.

Deniz, M., Schmitt Filho, A.L., Farley, J., de Quadros, S.F., Hötzel, M. (2019) High biodiversity silvopastoral system as an alternative to improve the thermal environment in the dairy farms. International journal of biometeorology 63, 83-92.

Derner, J.D., Boutton, T.W., Briske, D.D. (2006) Grazing and Ecosystem Carbon Storage in the North American Great Plains. Plant and Soil 280, 77-90.

Derner, J.D., Schuman, G.E. (2007) Carbon sequestration and rangelands: A synthesis of land management and precipitation effects. Journal of Soil and Water Conservation 62, 77.

Dillon, P.J.F. (2007) Achieving high dry-matter intake from pasture with grazing dairy cows. 1-26.

Dlamini, P., Chivenge, P., Manson, A., Chaplot, V. (2014) Land degradation impact on soil organic carbon and nitrogen stocks of sub-tropical humid grasslands in South Africa. Geoderma 235-236, 372-381.

Domingo, N.G.G., Balasubramanian, S., Thakrar, S.K., Clark, M.A., Adams, P.J., Marshall, J.D., Muller, N.Z., Pandis, S.N., Polasky, S., Robinson, A.L., Tessum, C.W., Tilman, D., Tschofen, P., Hill, J.D. (2021) Air quality–related health damages of food. Proceedings of the National Academy of Sciences 118, e2013637118.

Domke, G.M., Oswalt, S.N., Walters, B.F., Morin, R.S. (2020) Tree planting has the potential to increase carbon sequestration capacity of forests in the United States. Proceedings of the National Academy of Sciences 117, 24649.

Doran-Browne, N., Behrendt, R., Kingwell, R., Eckard, R. (2015) Modelling the potential of birdsfoot trefoil (*Lotus corniculatus*) to reduce methane emissions and increase production on wool and prime lamb farm enterprises. Animal Production Science 55, 1097-1105.

Doran-Browne, N., Wootton, M., Taylor, C., Eckard, R. (2018) Offsets required to reduce the carbon balance of sheep and beef farms through carbon sequestration in trees and soils. Animal Production Science 58, 1648-1655.

Doran-Browne, N.A., Ive, J., Graham, P., Eckard, R.J. (2016) Carbon-neutral wool farming in southeastern Australia. Animal Production Science 56, 417-422.

Douglas, G., Mackay, A., Vibart, R., Dodd, M., McIvor, I., McKenzie, C. (2020) Soil carbon stocks under grazed pasture and pasture-tree systems. Science of The Total Environment 715, 136910.

Eady, S., Carre, A., Grant, T. (2012) Life cycle assessment modelling of complex agricultural systems with multiple food and fibre co-products. Journal of Cleaner Production 28, 143-149.

Earles, J.M., Halog, A. (2011) Consequential life cycle assessment: a review. The International Journal of Life Cycle Assessment 16, 445-453.

FAO (2006a) Livestock's Long Shadow – Environmental Issues and Options. Food and Agriculture Organization of the United Nations (FAO), Rome, Italy.

FAO (2006b) Livestock's Long Shadow – Environmental Issues and Options. Food and Agriculture Organization of the United Nations (FAO), Rome, Italy (2006).

FAO (2019) Food and Agricultural Organization of the United Nations. Five practical actions towards low-carbon livestock. Available: <u>http://www.fao.org/3/ca7089en/ca7089en.pdf</u> [Accessed 18 July 2021].

FAO (2021a) Food and Agricultural Organization of the United Nations. Key facts and figures. Available:

http://www.fao.org/news/story/en/item/197623/icode/#:~:text=Total%20emissions%20from%20gl obal%20livestock,of%20all%20anthropogenic%20GHG%20emissions.&text=Cutting%20across%20all %20activities%20and,of%20the%20livestock%20sector's%20emissions. [Accessed 10 Jan 2021].

FAO (2021b) Food and Agriculture Organization of the United Nations. Global Livestock Environmental Assessment Model (GLEAM). Available: <u>http://www.fao.org/gleam/results/en/</u> [Accessed 18 July 2021].

FAOSTAT (2020) New Food Balances. <u>http://www.fao.org/faostat/en/#data/FBS</u>. [Accessed 28 March 2021].

Fleming, A., Stitzlein, C., Jakku, E., Fielke, S. (2019) Missed opportunity? Framing actions around cobenefits for carbon mitigation in Australian agriculture. Land Use Policy 85, 230-238.

Focus (2021) <u>https://focus2030.org/Focus-2030-and-the-Sustainable-Development-Goals-at-the-heart-of-our-work</u>.

Frank, S., Havlík, P., Soussana, J.-F., Levesque, A., Valin, H., Wollenberg, E., Kleinwechter, U., Fricko, O., Gusti, M., Herrero, M., Smith, P., Hasegawa, T., Kraxner, F., Obersteiner, M. (2017) Reducing greenhouse gas emissions in agriculture without compromising food security? Environmental Research Letters 12, 105004.

Franzel, S., Carsan, S., Lukuyu, B., Sinja, J., Wambugu, C. (2014) Fodder trees for improving livestock productivity and smallholder livelihoods in Africa. Current Opinion in Environmental Sustainability 6, 98-103.

Galford, G.L., Peña, O., Sullivan, A.K., Nash, J., Gurwick, N., Pirolli, G., Richards, M., White, J., Wollenberg, E. (2020) Agricultural development addresses food loss and waste while reducing greenhouse gas emissions. Science of The Total Environment 699, 134318.

Galloway, C., Conradie, B., Prozesky, H., Esler, K. (2018) Opportunities to improve sustainability on commercial pasture-based dairy farms by assessing environmental impact. Agricultural Systems 166, 1-9.

Garnett, T. (2009) Livestock-related greenhouse gas emissions: impacts and options for policy makers. Environmental Science & Policy 12, 491-503.

Garnett, T., Godde, C.M., Muller, A., Roos, E., Smith, P., de Boer, I., zu Ermgassen, E.K.H.J., Herrero, M., van Middelaar, C., Schader, C., van Zanten, H., (2017) Grazed and confused? Ruminating on cattle, grazing systems, methane, nitrous oxide, the soil carbon question - and what it means for

greenhouse gas emissions. Food Climate Research Network. Oxford Martin Programme on the Future of Food. Environmental Change Institute, University of Oxford.

Garrity, D.P. (2004) Agroforestry and the achievement of the Millennium Development Goals. Agroforestry Systems 61, 5-17.

Gerber, M., Span, R.J.P.I., Paris (2008) An analysis of available mathematical models for anaerobic digestion of organic substances for production of biogas.

Gerber, P., Steinfeld, H., Henderson, B., Mottet, A., Opio, C., Dijkman, J., Falcucci, A., Tempio, G. (2013) Tackling climate change through livestock: a global assessment of emissions and mitigation opportunities. Food and Agriculture Organization of the United Nations (FAO).

Godde, C.M., Garnett, T., Thornton, P.K., Ash, A.J., Herrero, M. (2018) Grazing systems expansion and intensification: Drivers, dynamics, and trade-offs. Global Food Security 16, 93-105.

Godde, C.M., Mason-D'Croz, D., Mayberry, D.E., Thornton, P.K., Herrero, M. (2021) Impacts of climate change on the livestock food supply chain; a review of the evidence. Global Food Security 28, 100488.

Godfray, H.C.J., Beddington, J.R., Crute, I.R., Haddad, L., Lawrence, D., Muir, J.F., Pretty, J., Robinson, S., Thomas, S.M., Toulmin, C. (2010) Food Security: The Challenge of Feeding 9 Billion People. Science 327, 812.

Goldemberg, J. (2007) Ethanol for a Sustainable Energy Future. Science 315, 808.

Gregory, N.G. (1995) The role of shelterbelts in protecting livestock: A review. New Zealand Journal of Agricultural Research 38, 423-450.

Grossi, G., Goglio, P., Vitali, A., Williams, A.G. (2019) Livestock and climate change: impact of livestock on climate and mitigation strategies. Animal Frontiers 9, 69-76.

Guo, L.B., Cowie, A.L., Montagu, K.D., Gifford, R.M. (2008) Carbon and nitrogen stocks in a native pasture and an adjacent 16-year-old Pinus radiata D. Don. plantation in Australia. Agriculture, Ecosystems & Environment 124, 205-218.

Haines, A., McMichael, A.J., Smith, K.R., Roberts, I., Woodcock, J., Markandya, A., Armstrong, B.G., Campbell-Lendrum, D., Dangour, A.D., Davies, M., Bruce, N., Tonne, C., Barrett, M., Wilkinson, P. (2009) Public health benefits of strategies to reduce greenhouse-gas emissions: overview and implications for policy makers. The Lancet 374, 2104-2114.

Hansen, B., Reich, P., Lake, S.P., Cavagnaro, T. (2010) Minimum width requirements for riparian zones to protect flowing waters and to conserve biodiversity: a review and recommendations. Report to the Office of Water, Department of Sustainability and Environment. Monash University, Australia, pp. 1-151.

Harrison, M.T., Christie, K.M., Rawnsley, R.P. (2017a) Assessing the reliability of dynamical and historical climate forecasts in simulating hindcast pasture growth rates. Animal Production Science 57, 1525-1535.

Harrison, M.T., Christie, K.M., Rawnsley, R.P., Eckard, R.J. (2014a) Modelling pasture management and livestock genotype interventions to improve whole-farm productivity and reduce greenhouse gas emissions intensities. Animal Production Science 54, 2018-2028.

Harrison, M.T., Cullen, B.R., Armstrong, D. (2017b) Management options for dairy farms under climate change: Effects of intensification, adaptation and simplification on pastures, milk production and profitability. Agricultural Systems 155, 19-32.

Harrison, M.T., Cullen, B.R., Rawnsley, R.P. (2016a) Modelling the sensitivity of agricultural systems to climate change and extreme climatic events. Agricultural Systems 148, 135-148.

Harrison, M.T., Cullen, B.R., Tomkins, N.W., McSweeney, C., Cohn, P., Eckard, R.J. (2016b) The concordance between greenhouse gas emissions, livestock production and profitability of extensive beef farming systems. Animal Production Science 56, 370-384.

Harrison, M.T., Evans, J.R., Dove, H., Moore, A.D. (2011a) Recovery dynamics of rainfed winter wheat after livestock grazing 1. Growth rates, grain yields, soil water use and water-use efficiency. Crop and Pasture Science 62, 947-959.

Harrison, M.T., Evans, J.R., Dove, H., Moore, A.D. (2011b) Recovery dynamics of rainfed winter wheat after livestock grazing 2. Light interception, radiation-use efficiency and dry-matter partitioning. Crop and Pasture Science 62, 960-971.

Harrison, M.T., Evans, J.R., Moore, A.D. (2012a) Using a mathematical framework to examine physiological changes in winter wheat after livestock grazing: 1. Model derivation and coefficient calibration. Field Crops Research 136, 116-126.

Harrison, M.T., Evans, J.R., Moore, A.D. (2012b) Using a mathematical framework to examine physiological changes in winter wheat after livestock grazing: 2. Model validation and effects of grazing management. Field Crops Research 136, 127-137.

Harrison, M.T., Jackson, T., Cullen, B.R., Rawnsley, R.P., Ho, C., Cummins, L., Eckard, R.J. (2014b) Increasing ewe genetic fecundity improves whole-farm production and reduces greenhouse gas emissions intensities: 1. Sheep production and emissions intensities. Agricultural Systems 131, 23-33.

Harrison, M.T., McSweeney, C., Tomkins, N.W., Eckard, R.J. (2015) Improving greenhouse gas emissions intensities of subtropical and tropical beef farming systems using Leucaena leucocephala. Agricultural Systems 136, 138-146.

Hasegawa, T., Fujimori, S., Shin, Y., Tanaka, A., Takahashi, K., Masui, T. (2015) Consequence of Climate Mitigation on the Risk of Hunger. Environmental Science & Technology 49, 7245-7253.

Havlík, P., Valin, H., Herrero, M., Obersteiner, M., Schmid, E., Rufino, M.C., Mosnier, A., Thornton, P.K., Böttcher, H., Conant, R.T., Frank, S., Fritz, S., Fuss, S., Kraxner, F., Notenbaert, A. (2014) Climate change mitigation through livestock system transitions. Proceedings of the National Academy of Sciences 111, 3709.

Häyhä, T., Lucas, P.L., van Vuuren, D.P., Cornell, S.E., Hoff, H. (2016) From Planetary Boundaries to national fair shares of the global safe operating space — How can the scales be bridged? Global Environmental Change 40, 60-72.

Henderson, B.B., Gerber, P.J., Hilinski, T.E., Falcucci, A., Ojima, D.S., Salvatore, M., Conant, R.T. (2015) Greenhouse gas mitigation potential of the world's grazing lands: Modeling soil carbon and nitrogen fluxes of mitigation practices. Agriculture, Ecosystems & Environment 207, 91-100.

Hernandez, R.R., Hoffacker, M.K., Murphy-Mariscal, M.L., Wu, G.C., Allen, M.F. (2015) Solar energy development impacts on land cover change and protected areas. Proceedings of the National Academy of Sciences 112, 13579.

Herrero, M., Grace, D., Njuki, J., Johnson, N., Enahoro, D., Silvestri, S., Rufino, M.C. (2013a) The roles of livestock in developing countries. Animal 7, 3-18.

Herrero, M., Havlík, P., Valin, H., Notenbaert, A., Rufino, M.C., Thornton, P.K., Blümmel, M., Weiss, F., Grace, D., Obersteiner, M. (2013b) Biomass use, production, feed efficiencies, and greenhouse gas emissions from global livestock systems. Proceedings of the National Academy of Sciences 110, 20888.

Herrero, M., Henderson, B., Havlík, P., Thornton, P.K., Conant, R.T., Smith, P., Wirsenius, S., Hristov, A.N., Gerber, P., Gill, M., Butterbach-Bahl, K., Valin, H., Garnett, T., Stehfest, E. (2016) Greenhouse gas mitigation potentials in the livestock sector. Nature Climate Change 6, 452-461.

Herrero, M., Thornton, P.K., Mason-D'Croz, D., Palmer, J., Bodirsky, B.L., Pradhan, P., Barrett, C.B., Benton, T.G., Hall, A., Pikaar, I., Bogard, J.R., Bonnett, G.D., Bryan, B.A., Campbell, B.M., Christensen, S., Clark, M., Fanzo, J., Godde, C.M., Jarvis, A., Loboguerrero, A.M., Mathys, A., McIntyre, C.L., Naylor, R.L., Nelson, R., Obersteiner, M., Parodi, A., Popp, A., Ricketts, K., Smith, P., Valin, H., Vermeulen, S.J., Vervoort, J., van Wijk, M., van Zanten, H.H.E., West, P.C., Wood, S.A., Rockström, J. (2021) Articulating the effect of food systems innovation on the Sustainable Development Goals. The Lancet Planetary Health 5, e50-e62.

Herrero, M., Wirsenius, S., Henderson, B., Rigolot, C., Thornton, P., Havlík, P., de Boer, I., Gerber, P.J. (2015) Livestock and the Environment: What Have We Learned in the Past Decade? Annual Review of Environment and Resources 40, 177-202.

Herrick, J.E., Neff, J., Quandt, A., Salley, S., Maynard, J., Ganguli, A., Bestelmeyer, B. (2019) Prioritizing land for investments based on short- and long-term land potential and degradation risk: A strategic approach. Environmental Science & Policy 96, 52-58.

Ho, C.K.M., Jackson, T., Harrison, M.T., Eckard, R.J. (2014) Increasing ewe genetic fecundity improves whole-farm production and reduces greenhouse gas emissions intensities: 2. Economic performance. Animal Production Science 54, 1248-1253.

Honda, E.A., Durigan, G. (2016) Woody encroachment and its consequences on hydrological processes in the savannah. Philosophical Transactions of the Royal Society B: Biological Sciences 371, 20150313.

Hristov, A.N., Oh, J., Firkins, J.L., Dijkstra, J., Kebreab, E., Waghorn, G., Makkar, H.P.S., Adesogan, A.T., Yang, W., Lee, C., Gerber, P.J., Henderson, B., Tricarico, J.M. (2013) SPECIAL TOPICS — Mitigation of methane and nitrous oxide emissions from animal operations: I. A review of enteric methane mitigation options1. Journal of Animal Science 91, 5045-5069.

Hu, X., Kerckhof, F.-M., Ghesquière, J., Bernaerts, K., Boeckx, P., Clauwaert, P., Boon, N. (2020) Microbial Protein out of Thin Air: Fixation of Nitrogen Gas by an Autotrophic Hydrogen-Oxidizing Bacterial Enrichment. Environmental Science & Technology 54, 3609-3617.

IPCC (2014) Contribution of Working Group III to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change (Cambridge, NY: Cambridge University Press) Climate change 2014: mitigation of climate change. 1454.

Jerónimo, E., Pinheiro, C., Lamy, E., Dentinho, M.T., Sales-Baptista, E., Lopes, O., Silva, F. (2016) Tannins in ruminant nutrition: Impact on animal performance and quality of edible products.

Jiang, Z.-Y., Hu, Z.-M., Lai, D.Y.F., Han, D.-R., Wang, M., Liu, M., Zhang, M., Guo, M.-Y. (2020) Light grazing facilitates carbon accumulation in subsoil in Chinese grasslands: A meta-analysis. Global Change Biology 26, 7186-7197.

Jindal, R., Swallow, B., Kerr, J. (2008) Forestry-based carbon sequestration projects in Africa: Potential benefits and challenges. Natural Resources Forum 32, 116-130.

Jones, I.J., MacDonald, A.J., Hopkins, S.R., Lund, A.J., Liu, Z.Y.-C., Fawzi, N.I., Purba, M.P., Fankhauser, K., Chamberlin, A.J., Nirmala, M., Blundell, A.G., Emerson, A., Jennings, J., Gaffikin, L., Barry, M., Lopez-Carr, D., Webb, K., De Leo, G.A., Sokolow, S.H. (2020) Improving rural health care reduces illegal logging and conserves carbon in a tropical forest. Proceedings of the National Academy of Sciences 117, 28515.

Joseph, S., Pow, D., Dawson, K., Mitchell, D.R.G., Rawal, A., Hook, J., Taherymoosavi, S., Van Zwieten, L., Rust, J., Donne, S., Munroe, P., Pace, B., Graber, E., Thomas, T., Nielsen, S., Ye, J., Lin, Y., Pan, G., Li, L., Solaiman, Z.M. (2015) Feeding Biochar to Cows: An Innovative Solution for Improving Soil Fertility and Farm Productivity. Pedosphere 25, 666-679.

Kemp, D.R., Guodong, H., Xiangyang, H., Michalk, D.L., Fujiang, H., Jianping, W., Yingjun, Z. (2013) Innovative grassland management systems for environmental and livelihood benefits. Proceedings of the National Academy of Sciences 110, 8369-8374.

Kinley, R.D., Martinez-Fernandez, G., Matthews, M.K., de Nys, R., Magnusson, M., Tomkins, N.W. (2020) Mitigating the carbon footprint and improving productivity of ruminant livestock agriculture using a red seaweed. Journal of Cleaner Production 259, 120836.

Kitalyi, A., Mtenga, L., Morton, J., McLeod, A., Thornton, P., Dorward, A., Saadhullah, M., Owen, E., Jayasuriya, A., Smith, T. (2005) Why keep livestock if you are poor? In Livestock and wealth creation: improving the husbandry of animals kept by resource-poor people in developing countries (eds, Owen E., Kitalyi A., Jayasuriay N.& Smith T.), pp. 13–27. Nottingham, UK: Nottingham University Press. .

Klerkx, L., Begemann, S. (2020) Supporting food systems transformation: The what, why, who, where and how of mission-oriented agricultural innovation systems. Agricultural Systems 184, 102901.

Kongsager, R., Napier, J., Mertz, O. (2013) The carbon sequestration potential of tree crop plantations. Mitigation and Adaptation Strategies for Global Change 18, 1197-1213.

Kösler, J.E., Calvo, O.C., Franzaring, J., Fangmeier, A. (2019) Evaluating the ecotoxicity of nitrification inhibitors using terrestrial and aquatic test organisms. Environmental Sciences Europe 31, 91.

Kragt, M.E., Dumbrell, N.P., Blackmore, L. (2017) Motivations and barriers for Western Australian broad-acre farmers to adopt carbon farming. Environmental Science & Policy 73, 115-123.

Kreidenweis, U., Humpenöder, F., Stevanović, M., Bodirsky, B.L., Kriegler, E., Lotze-Campen, H., Popp, A. (2016) Afforestation to mitigate climate change: impacts on food prices under consideration of albedo effects. Environmental Research Letters 11, 085001.

Kristjanson, P., Waters-Bayer, A., Johnson, N., Tipilda, A., Njuki, J., Baltenweck, I., Grace, D., MacMillan, S., (2014) Livestock and Women's Livelihoods, in: Quisumbing, A.R., Meinzen-Dick, R., Raney, T.L., Croppenstedt, A., Behrman, J.A., Peterman, A. (Eds.), Gender in Agriculture: Closing the Knowledge Gap. Springer Netherlands, Dordrecht, pp. 209-233.

Kumari, S., Fagodiya, R.K., Hiloidhari, M., Dahiya, R.P., Kumar, A. (2020) Methane production and estimation from livestock husbandry: A mechanistic understanding and emerging mitigation options. Science of The Total Environment 709, 136135.

Laborde, D., Mamun, A., Martin, W., Piñeiro, V., Vos, R. (2021) Agricultural subsidies and global greenhouse gas emissions. Nature Communications 12, 2601.

Lade, S.J., Steffen, W., de Vries, W., Carpenter, S.R., Donges, J.F., Gerten, D., Hoff, H., Newbold, T., Richardson, K., Rockström, J. (2020) Human impacts on planetary boundaries amplified by Earth system interactions. Nature Sustainability 3, 119-128.

Lal, R. (2004) Soil Carbon Sequestration Impacts on Global Climate Change and Food Security. Science 304, 1623.

Langworthy, A.D., Rawnsley, R.P., Freeman, M.J., Pembleton, K.G., Corkrey, R., Harrison, M.T., Lane, P.A., Henry, D.A. (2018) Potential of summer-active temperate (C3) perennial forages to mitigate the detrimental effects of supraoptimal temperatures on summer home-grown feed production in south-eastern Australian dairying regions. Crop and Pasture Science 69, 808-820.

Larson, Z., (2000) Selections from nonviolence and culture: finding the Dharma in a flesh-based diet. Excerpted from: Nonviolence in a Tibetan culture: a glimpse at how Tibetans view and practice nonviolence in politics and daily life. Madison, WI. University of Wisconsin.

Lawrence, A.B., Conington, J., Simm, G. (2004) Breeding and animal welfare: practical and theoretical advantages of multi-trait selection. Animal Welfare 13, 191-196.

Leahy, S., Clark, H., Reisinger, A., (2020) Challenges and Prospects for Agricultural Greenhouse Gas Mitigation Pathways Consistent With the Paris Agreement, Frontiers in Sustainable Food Systems.

Leahy, S.C., Kearney, L., Reisinger, A., Clark, H. (2019) Mitigating greenhouse gas emissions from New Zealand pasture-based livestock farm systems. Journal of New Zealand Grasslands, 101-110.

Lee, J., Ingalls, M., Erickson, J.D., Wollenberg, E. (2016) Bridging organizations in agricultural carbon markets and poverty alleviation: An analysis of pro-Poor carbon market projects in East Africa. Global Environmental Change 39, 98-107.

Leip, A., Billen, G., Garnier, J., Grizzetti, B., Lassaletta, L., Reis, S., Simpson, D., Sutton, M.A., de Vries, W., Weiss, F., Westhoek, H. (2015) Impacts of European livestock production: nitrogen, sulphur, phosphorus and greenhouse gas emissions, land-use, water eutrophication and biodiversity. Environmental Research Letters 10, 115004.

Lira Junior, M.A., Fracetto, F.J.C., Ferreira, J.d.S., Silva, M.B., Fracetto, G.G.M. (2020) Legume-based silvopastoral systems drive C and N soil stocks in a subhumid tropical environment. CATENA 189, 104508.

Liu, N., Zhang, Y., Chang, S., Kan, H., Lin, L. (2012) Impact of Grazing on Soil Carbon and Microbial Biomass in Typical Steppe and Desert Steppe of Inner Mongolia. PLOS ONE 7, e36434.

Lohbeck, M., Winowiecki, L., Aynekulu, E., Okia, C., Vågen, T.-G. (2018) Trait-based approaches for guiding the restoration of degraded agricultural landscapes in East Africa. Journal of Applied Ecology 55, 59-68.

Lonsdale, W.M. (1994) Inviting trouble: Introduced pasture species in northern Australia. Australian Journal of Ecology 19, 345-354.

Lovarelli, D., Finzi, A., Mattachini, G., Riva, E. (2020) A Survey of Dairy Cattle Behavior in Different Barns in Northern Italy. Animals 10.

Ludemann, C.I., Howden, S.M., Eckard, R.J. (2016) What is the best use of oil from cotton (<i>Gossypium</i> spp.) and canola (<i>Brassica</i> spp.) for reducing net greenhouse gas emissions: biodiesel, or as a feed for cattle? Animal Production Science 56, 442-450.

Luo, J., Balvert, S.F., Wise, B., Welten, B., Ledgard, S.F., de Klein, C.A.M., Lindsey, S., Judge, A. (2018) Using alternative forage species to reduce emissions of the greenhouse gas nitrous oxide from cattle urine deposited onto soil. Science of The Total Environment 610-611, 1271-1280.

Lynch, J., Cain, M., Pierrehumbert, R., Allen, M. (2020) Demonstrating GWP: a means of reporting warming-equivalent emissions that captures the contrasting impacts of short- and long-lived climate pollutants. Environmental Research Letters 15.

Lynch, J., Pierrehumbert, R. (2019) Climate Impacts of Cultured Meat and Beef Cattle. Frontiers in Sustainable Food Systems 3.

MacLeod, M., Moran, D. (2017) Integrating livestock health measures into marginal abatement cost curves. Revue Scientifique et Technique-Office International des Épizooties 36, 97-104.

Manning, A.D., Fischer, J., Lindenmayer, D.B. (2006) Scattered trees are keystone structures – Implications for conservation. Biological Conservation 132, 311-321.

Martin, C., Morgavi, D.P., Doreau, M. (2010) Methane mitigation in ruminants: from microbe to the farm scale. Animal 4, 351-365.

Martínez, J., Cajas, Y.S., León, J.D., Osorio, N.W. (2014) Silvopastoral Systems Enhance Soil Quality in Grasslands of Colombia. Applied and Environmental Soil Science 2014, 359736.

Mayberry, D., Bartlett, H., Moss, J., Davison, T., Herrero, M. (2019) Pathways to carbon-neutrality for the Australian red meat sector. Agricultural Systems 175, 13-21.

Mayer, F.D., Brondani, M., Vasquez Carrillo, M.C., Hoffmann, R., Silva Lora, E.E. (2020) Revisiting energy efficiency, renewability, and sustainability indicators in biofuels life cycle: Analysis and standardization proposal. Journal of Cleaner Production 252, 119850.

Meier, E.A., Thorburn, P.J., Bell, L.W., Harrison, M.T., Biggs, J.S. (2020) Greenhouse Gas Emissions From Cropping and Grazed Pastures Are Similar: A Simulation Analysis in Australia. Frontiers in Sustainable Food Systems 3, 121.

Menzi, H., Oenema, O., Burton, C., Shipin, O., Gerber, P., Robinson, T., Franceschini, G. (2010) Impacts of intensive livestock production and manure management on the environment. Livestock in a changing landscape 1, 139-163. Meyer, R., Cullen, B.R., Eckard, R.J. (2016) Modelling the influence of soil carbon on net greenhouse gas emissions from grazed pastures. Animal Production Science 56, 585-593.

Meyer, R., Cullen, B.R., Johnson, I.R., Eckard, R.J. (2015) Process modelling to assess the sequestration and productivity benefits of soil carbon for pasture. Agriculture, Ecosystems & Environment 213, 272-280.

Michalk, D.L., Kemp, D.R., Badgery, W.B., Wu, J., Zhang, Y., Thomassin, P.J. (2019) Sustainability and future food security—A global perspective for livestock production. Land Degradation & Development 30, 561-573.

MLA (2016) <u>https://www.mla.com.au/globalassets/mla-corporate/blocks/research-and-development/erf-fact-sheet_nitrates_final.pdf</u>

MLA (2020) <u>https://www.mla.com.au/globalassets/mla-corporate/blocks/research-and-development/tt_bmp-for-feeding-nitrates-to-cattle_web.pdf</u>.

Modernel, P., Astigarraga, L., Picasso, V. (2013) Global versus local environmental impacts of grazing and confined beef production systems. Environmental Research Letters 8, 035052.

Molossi, L., Hoshide, A.K., Pedrosa, L.M., Oliveira, A.S., Abreu, D.C. (2020) Improve Pasture or Feed Grain? Greenhouse Gas Emissions, Profitability, and Resource Use for Nelore Beef Cattle in Brazil's Cerrado and Amazon Biomes. Animals 10.

Montagnini, F., Ibrahim, M., Murgueitio, E.J.B.e.f.d.t. (2013) Silvopastoral systems and climate change mitigation in Latin America. 316, 3-16.

Mottet, A., de Haan, C., Falcucci, A., Tempio, G., Opio, C., Gerber, P. (2017) Livestock: On our plates or eating at our table? A new analysis of the feed/food debate. Global Food Security 14, 1-8.

Myhre, G., Shindell, D., Bréon, F.-M., Collins, W., Fuglestvedt, J., Huang, J., Koch, D., Lamarque, J.-F., Lee, D., Mendoza, B., Nakajima, T., Robock, A., Stephens, G., T., T., Zhang, H., (2013) Anthropogenic and Natural Radiative Forcing. In: Climate Change 2013: The Physical Science Basis, in: Stocker, T.F., Qin, D., Plattner, G.-K., Tignor, M., Allen, S.K., Boschung, J., Nauels, A., Xia, Y., V., B., P.M., M. (Eds.), Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA.

Nakyinsige, K., Man, Y.B.C., Sazili, A.Q. (2012) Halal authenticity issues in meat and meat products. Meat Science 91, 207-214.

Neumann, C.G., Bwibo, N.O., Murphy, S.P., Sigman, M., Whaley, S., Allen, L.H., Guthrie, D., Weiss, R.E., Demment, M.W. (2003) Animal Source Foods Improve Dietary Quality, Micronutrient Status, Growth and Cognitive Function in Kenyan School Children: Background, Study Design and Baseline Findings. The Journal of Nutrition 133, 3941S-3949S.

Nigussie, Z., Tsunekawa, A., Haregeweyn, N., Tsubo, M., Adgo, E., Ayalew, Z., Abele, S. (2021) The impacts of Acacia decurrens plantations on livelihoods in rural Ethiopia. Land Use Policy 100, 104928.

Nilsson, S., Schopfhauser, W. (1995) The carbon-sequestration potential of a global afforestation program. Climatic Change 30, 267-293.

NRC (2013) Sustainable development of algal biofuels in the United States. Chapter: 5 Environmental Effects. National Academies Press.

O'Connell, D., Maru, Y., Grigg, N., Walker, B., Abel, N., Wise, R., Cowie, A., Butler, J., Stone-Jovicich, S., STAFFORD SMITH, M., (2019) Resilience, Adaptation Pathways and Transformation Approach: A guide for designing, implementing and assessing interventions for sustainable futures (version 2). <u>https://acfid.asn.au/sites/site.acfid/files/19-00418_LW_REPORT_RAPTAGuide_WEB_190829.pdf</u>. Australia.

OECD (2021) OECD Agriculture Statistics. Available: <u>https://www.oecd-ilibrary.org/agriculture-and-food/data/oecd-agriculture-statistics_agr-data-en</u> [Accessed 14 July 2021].

Orgill, S., Condon, J., Conyers, M., Greene, R., Murphy, B., (2012) Opportunities to sequester carbon in soil: management of perennial pastures, Capturing Opportunities and Overcoming Obstacles in Australian Agronomy. Proceedings of the 16th Australian Agronomy Conference'. Armidale, NSW.(Ed. I. Yunusa.)(Australian Society of Agronomy/The Regional Institute Inc.: Gosford, NSW.).

Palmer, J., Thorburn, P.J., Biggs, J.S., Dominati, E.J., Probert, M.E., Meier, E.A., Huth, N.I., Dodd, M., Snow, V., Larsen, J.R., Parton, W.J. (2017) Nitrogen Cycling from Increased Soil Organic Carbon Contributes Both Positively and Negatively to Ecosystem Services in Wheat Agro-Ecosystems. Frontiers in Plant Science 8.

Pang, D., Cui, M., Liu, Y., Wang, G., Cao, J., Wang, X., Dan, X., Zhou, J. (2019) Responses of soil labile organic carbon fractions and stocks to different vegetation restoration strategies in degraded karst ecosystems of southwest China. Ecological Engineering 138, 391-402.

Paul, K.I., Roxburgh, S.H., Chave, J., England, J.R., Zerihun, A., Specht, A., Lewis, T., Bennett, L.T., Baker, T.G., Adams, M.A., Huxtable, D., Montagu, K.D., Falster, D.S., Feller, M., Sochacki, S., Ritson, P., Bastin, G., Bartle, J., Wildy, D., Hobbs, T., Larmour, J., Waterworth, R., Stewart, H.T.L., Jonson, J., Forrester, D.I., Applegate, G., Mendham, D., Bradford, M., O'Grady, A., Green, D., Sudmeyer, R., Rance, S.J., Turner, J., Barton, C., Wenk, E.H., Grove, T., Attiwill, P.M., Pinkard, E., Butler, D., Brooksbank, K., Spencer, B., Snowdon, P., O'Brien, N., Battaglia, M., Cameron, D.M., Hamilton, S., McAuthur, G., Sinclair, J. (2016) Testing the generality of above-ground biomass allometry across plant functional types at the continent scale. Global Change Biology 22, 2106-2124.

Pembleton, K.G., Cullen, B.R., Rawnsley, R.P., Harrison, M.T., Ramilan, T. (2016) Modelling the resilience of forage crop production to future climate change in the dairy regions of Southeastern Australia using APSIM. The Journal of Agricultural Science 154, 1131-1152.

Petersen, S.O., Blanchard, M., Chadwick, D., Del Prado, A., Edouard, N., Mosquera, J., Sommer, S.G. (2013) Manure management for greenhouse gas mitigation. Animal 7, 266-282.

Phelan, D.C., Harrison, M.T., Kemmerer, E.P., Parsons, D. (2015) Management opportunities for boosting productivity of cool-temperate dairy farms under climate change. Agricultural Systems 138, 46-54.

Pinto, M., Merino, P., del Prado, A., Estavillo, J.M., Yamulki, S., Gebauer, G., Piertzak, S., Lauf, J., Oenema, O. (2004) Increased emissions of nitric oxide and nitrous oxide following tillage of a perennial pasture. Nutrient Cycling in Agroecosystems 70, 13-22.

Plevin, R.J., Delucchi, M.A., Creutzig, F. (2014) Using Attributional Life Cycle Assessment to Estimate Climate-Change Mitigation Benefits Misleads Policy Makers. Journal of Industrial Ecology 18, 73-83.

Poore, J., Nemecek, T. (2018) Reducing food's environmental impacts through producers and consumers. Science 360, 987.

Popp, A., Calvin, K., Fujimori, S., Havlik, P., Humpenöder, F., Stehfest, E., Bodirsky, B.L., Dietrich, J.P., Doelmann, J.C., Gusti, M., Hasegawa, T., Kyle, P., Obersteiner, M., Tabeau, A., Takahashi, K., Valin, H., Waldhoff, S., Weindl, I., Wise, M., Kriegler, E., Lotze-Campen, H., Fricko, O., Riahi, K., Vuuren, D.P.v. (2017) Land-use futures in the shared socio-economic pathways. Global Environmental Change 42, 331-345.

Popp, A., Lotze-Campen, H., Bodirsky, B. (2010) Food consumption, diet shifts and associated non-CO2 greenhouse gases from agricultural production. Global Environmental Change 20, 451-462.

Radrizzani, A., Shelton, H.M., Dalzell, S.A., Kirchhof, G. (2011) Soil organic carbon and total nitrogen under *Leucaena leucocephala* pastures in Queensland. Crop and Pasture Science 62, 337-345.

Rae, A., Nayga, R. (2010) Trends in consumption, production and trade in livestock and livestock products. Island Press.

Ramachandran Nair, P.K., Nair, V.D., Mohan Kumar, B., Showalter, J.M., (2010) Chapter Five - Carbon Sequestration in Agroforestry Systems, in: Sparks, D.L. (Ed.), Advances in Agronomy. Academic Press, pp. 237-307.

Ramprasad, V., Joglekar, A., Fleischman, F. (2020) Plantations and pastoralists: afforestation activities make pastoralists in the Indian Himalaya vulnerable. Ecology and Society 25.

Rawnsley, R., Dynes, R.A., Christie, K.M., Harrison, M.T., Doran-Browne, N.A., Vibart, R., Eckard, R. (2018) A review of whole farm-system analysis in evaluating greenhouse-gas mitigation strategies from livestock production systems. Animal Production Science 58, 980-989.

Reijnders, L. (2012) Life cycle assessment of greenhouse gas emissions. Springer reference.

Reisinger, A., Clark, H. (2018) How much do direct livestock emissions actually contribute to global warming? Global Change Biology 24, 1749-1761.

Rice, P., O'Brien, D., Shalloo, L., Holden, N.M. (2017) Evaluation of allocation methods for calculation of carbon footprint of grass-based dairy production. Journal of Environmental Management 202, 311-319.

Ritchie, H., (2019) Half of the world's habitable land is used for agriculture. Our World in Data. Global land use for food production. Available: <u>https://ourworldindata.org/global-land-for-agriculture</u> [Accessed 29 March 2021].

Ritchie, H., Roser, M. (2020) Environmental impacts of food production. Our world in data.

Rivera-Ferre, M.G., López-i-Gelats, F., Howden, M., Smith, P., Morton, J.F., Herrero, M. (2016) Reframing the climate change debate in the livestock sector: mitigation and adaptation options. WIREs Climate Change 7, 869-892.

Roberts, K.G., Gloy, B.A., Joseph, S., Scott, N.R., Lehmann, J. (2010) Life Cycle Assessment of Biochar Systems: Estimating the Energetic, Economic, and Climate Change Potential. Environmental Science & Technology 44, 827-833.

Rockström, J., Steffen, W., Noone, K., Persson, Å., Chapin Iii, F.S., Lambin, E., Lenton, T.M., Scheffer, M., Folke, C., Schellnhuber, H. (2009) Planetary boundaries: Exploring the safe operating space for humanity. Ecology and Society 14, 32.

Rogelj, J., Schleussner, C.-F. (2019) Unintentional unfairness when applying new greenhouse gas emissions metrics at country level. Environmental Research Letters 14, 114039.

Rohman, A., Che Man, Y.B. (2012) Analysis of Pig Derivatives for Halal Authentication Studies. Food Reviews International 28, 97-112.

Rolfe, J., (2001) Economics of reducing methane emissions from cattle production in central Queensland. Meat & Livestock Australia. Available: <u>https://www.mla.com.au/contentassets/b0a03a4317e446f2a939dbbff6b96ab3/nap3.226_final_rep_ort.pdf</u> [Accessed 29 March 2021].

Roque, B.M., Brooke, C.G., Ladau, J., Polley, T., Marsh, L.J., Najafi, N., Pandey, P., Singh, L., Kinley, R., Salwen, J.K., Eloe-Fadrosh, E., Kebreab, E., Hess, M. (2019) Effect of the macroalgae Asparagopsis taxiformis on methane production and rumen microbiome assemblage. Animal Microbiome 1, 3.

Rose, S., Lee, H.-L. (2008) Non-CO₂ greenhouse gas emissions data for climate change economic analysis. GTAP working paper No. 43.

Rubin, D., Tezera, S., Caldwell, L. (2010) A calf, a house, a business of one's own: Microcredit, asset accumulation, and economic empowerment in GL CRSP projects in Ethiopia and Ghana. Global livestock collaborative research support program.

Ruis, S.J., Blanco-Canqui, H. (2017) Cover Crops Could Offset Crop Residue Removal Effects on Soil Carbon and Other Properties: A Review. Agronomy Journal 109, 1785-1805.

Salmon, G., Teufel, N., Baltenweck, I., van Wijk, M., Claessens, L., Marshall, K. (2018) Trade-offs in livestock development at farm level: Different actors with different objectives. Global Food Security 17, 103-112.

Sanderson, J.S., Beutler, C., Brown, J.R., Burke, I., Chapman, T., Conant, R.T., Derner, J.D., Easter, M., Fuhlendorf, S.D., Grissom, G., Herrick, J.E., Liptzin, D., Morgan, J.A., Murph, R., Pague, C., Rangwala, I., Ray, D., Rondeau, R., Schulz, T., Sullivan, T. (2020) Cattle, conservation, and carbon in the western Great Plains. Journal of Soil and Water Conservation 75, 5A.

Sándor, R., Ehrhardt, F., Grace, P., Recous, S., Smith, P., Snow, V., Soussana, J.-F., Basso, B., Bhatia, A., Brilli, L., Doltra, J., Dorich, C.D., Doro, L., Fitton, N., Grant, B., Harrison, M.T., Kirschbaum, M.U.F., Klumpp, K., Laville, P., Léonard, J., Martin, R., Massad, R.-S., Moore, A., Myrgiotis, V., Pattey, E., Rolinski, S., Sharp, J., Skiba, U., Smith, W., Wu, L., Zhang, Q., Bellocchi, G. (2020) Ensemble modelling of carbon fluxes in grasslands and croplands. Field Crops Research 252, 107791.

Schellnhuber, H.J., Cramer, W., Nakicenovic, N., Wigley, T., Yohe, G. (2006) Avoiding Dangerous Climate Change, Cambridge University Press.

Schleussner, C.F., Lissner, T.K., Fischer, E.M., Wohland, J., Perrette, M., Golly, A., Rogelj, J., Childers, K., Schewe, J., Frieler, K., Mengel, M., Hare, W., Schaeffer, M. (2016) Differential climate impacts for policy-relevant limits to global warming: the case of 1.5 °C and 2 °C. Earth Syst. Dynam. 7, 327-351.

Schmidt, H.-P., Hagemann, N., Draper, K., Kammann, C. (2019) The use of biochar in animal feeding. PeerJ 7, e7373. Scurlock, J.M.O., Hall, D.O. (1998) The global carbon sink: a grassland perspective. Global Change Biology 4, 229-233.

Shields, S., Orme-Evans, G. (2015) The Impacts of Climate Change Mitigation Strategies on Animal Welfare. Animals 5.

Shine, K.P. (2009) The global warming potential—the need for an interdisciplinary retrial. Climatic Change 96, 467-472.

Shine, K.P., Berntsen, T.K., Fuglestvedt, J.S., Skeie, R.B., Stuber, N. (2007) Comparing the climate effect of emissions of short- and long-lived climate agents. Philosophical Transactions of the Royal Society of London A: Mathematical, Physical and Engineering Sciences 365, 1903-1914.

Smetana, S., Mathys, A., Knoch, A., Heinz, V. (2015) Meat alternatives: life cycle assessment of most known meat substitutes. The International Journal of Life Cycle Assessment 20, 1254-1267.

Smith, A.P., Christie, K.M., Harrison, M.T., Eckard, R.J. (2021) Ammonia volatilisation from grazed, pasture based dairy farming systems. Agricultural Systems 190, 103119.

Smith, P. (2012) Agricultural greenhouse gas mitigation potential globally, in Europe and in the UK: what have we learnt in the last 20 years? Global Change Biology 18, 35-43.

Smith, P., Calvin, K., Nkem, J., Campbell, D., Cherubini, F., Grassi, G., Korotkov, V., Le Hoang, A., Lwasa, S., McElwee, P., Nkonya, E., Saigusa, N., Soussana, J.-F., Taboada, M.A., Manning, F.C., Nampanzira, D., Arias-Navarro, C., Vizzarri, M., House, J., Roe, S., Cowie, A., Rounsevell, M., Arneth, A. (2020) Which practices co-deliver food security, climate change mitigation and adaptation, and combat land degradation and desertification? Global Change Biology 26, 1532-1575.

Smith, P., Haberl, H., Popp, A., Erb, K.-h., Lauk, C., Harper, R., Tubiello, F., de Siqueira Pinto, A., Jafari, M., Sohi, S., Masera, O., Böttcher, H., Berndes, G., Bustamante, M., Ahammad, H., Clark, H., Dong, H., Elsiddig, E., Mbow, C., Ravindranath, N., Rice, C., Robledo Abad, C., Romanovskaya, A., Sperling, F., Herrero, M., House, J., Rose, S. (2013) How much land-based greenhouse gas mitigation can be achieved without compromising food security and environmental goals? Global Change Biology 19, 2285-2302.

Smith, P., Martino, D., Cai, Z., Gwary, D., Janzen, H., Kumar, P., McCarl, B., Ogle, S., O'Mara, F., Rice, C., Scholes, B., Sirotenko, O., Howden, M., McAllister, T., Pan, G., Romanenkov, V., Schneider, U., Towprayoon, S. (2007) Policy and technological constraints to implementation of greenhouse gas mitigation options in agriculture. Agriculture, Ecosystems & Environment 118, 6-28.

Spiller, M., Muys, M., Papini, G., Sakarika, M., Buyle, M., Vlaeminck, S.E. (2020) Environmental impact of microbial protein from potato wastewater as feed ingredient: Comparative consequential life cycle assessment of three production systems and soybean meal. Water Research 171, 115406.

Steffen, W., Richardson, K., Rockström, J., Cornell, S.E., Fetzer, I., Bennett, E.M., Biggs, R., Carpenter, S.R., De Vries, W., De Wit, C.A.J.S. (2015) Planetary boundaries: Guiding human development on a changing planet. Science 347.

Stehfest, E., Berg, M.v.d., Woltjer, G., Msangi, S., Westhoek, H. (2013) Options to reduce the environmental effects of livestock production – Comparison of two economic models. Agricultural Systems 114, 38-53.

Stehfest, E., Bouwman, L., van Vuuren, D.P., den Elzen, M.G.J., Eickhout, B., Kabat, P. (2009) Climate benefits of changing diet. Climatic Change 95, 83-102.

Steinfeld, H., Gerber, P. (2010) Livestock production and the global environment: Consume less or produce better? Proceedings of the National Academy of Sciences 107, 18237-18238.

Steinfeld, H., Gerber, P., Wassenaar, T., Castel, V., Rosales, M., de Haan, C. (2006a) Livestock's long shadow: environmental issues and options. Food and Agriculture Organization, Rome, Italy.

Steinfeld, H., Gerber, P., Wassenaar, T., Castel, V., Rosales, M., Rosales, M., de Haan, C. (2006b) Livestock's long shadow: environmental issues and options. Food & Agriculture Org.

Steinfeld, H., Haan, C.d., Blackburn, H. (2003) Livestock – Environment Interactions. Issues and options. Food and Agricultural Organisation of the United Nations. Chapter 1: livestock, environment and human needs. Available:

http://www.fao.org/3/x5305e/x5305e02.htm#livestock%20in%20a%20changing%20world [Accessed 10 January 2021].

Struik, P.C., Kuyper, T.W., Brussaard, L., Leeuwis, C. (2014) Deconstructing and unpacking scientific controversies in intensification and sustainability: why the tensions in concepts and values? Current Opinion in Environmental Sustainability 8, 80-88.

Sun, X., Pacheco, D., Luo, D. (2016) Forage brassica: a feed to mitigate enteric methane emissions? Animal Production Science 56, 451-456.

Suybeng, B., Charmley, E., Gardiner, C.P., Malau-Aduli, B.S., Malau-Aduli, A.E.O. (2019) Methane Emissions and the Use of Desmanthus in Beef Cattle Production in Northern Australia. Animals 9.

Taylor, C., Eckard, R. (2016) Comparative analysis of greenhouse gas emissions from three beef cattle herds in a corporate farming enterprise. Animal Production Science 56, 482-494.

Taylor, C.A., Harrison, M.T., Telfer, M., Eckard, R. (2016) Modelled greenhouse gas emissions from beef cattle grazing irrigated leucaena in northern Australia. Animal Production Science 56, 594-604.

Tengberg, A., Valencia, S. (2018) Integrated approaches to natural resources management—Theory and practice. Land Degradation & Development 29, 1845-1857.

Tessema, B., Sommer, R., Piikki, K., Söderström, M., Namirembe, S., Notenbaert, A., Tamene, L., Nyawira, S., Paul, B. (2020) Potential for soil organic carbon sequestration in grasslands in East African countries: A review. Grassland Science 66, 135-144.

Thomson, B.C., Hammond, K.J., Muir, P.D. (2016) A review of greenhouse gas emissions from the use of brassica and fodder beet forages on New Zealand farms. Prepared for NZAGRC/PGGRC. Available: <u>https://www.nzagrc.org.nz/user/file/1752/OFR_Brassica_Final_Report_June%202016_web%20copy.pdf</u>

Thornton, P.K. (2010) Livestock production: recent trends, future prospects. Philosophical Transactions of the Royal Society B: Biological Sciences 365, 2853-2867.

Thornton, P.K., Gerber, P.J. (2010) Climate change and the growth of the livestock sector in developing countries. Mitigation and Adaptation Strategies for Global Change 15, 169-184.

Thornton, P.K., Whitbread, A., Baedeker, T., Cairns, J., Claessens, L., Baethgen, W., Bunn, C., Friedmann, M., Giller, K.E., Herrero, M., Howden, M., Kilcline, K., Nangia, V., Ramirez-Villegas, J., Kumar, S., West, P.C., Keating, B. (2018) A framework for priority-setting in climate smart agriculture research. Agricultural Systems 167, 161-175.

Tilman, D., Clark, M. (2014) Global diets link environmental sustainability and human health. Nature 515, 518-522.

Tomkins, N., Harrison, M., McSweeney, C.S., Denman, S., Charmley, E., Lambrides, C.J., Dalal, R. (2019) Greenhouse gas implications of leucaena-based pastures. Can we develop an emissions reduction methodology for the beef industry? Tropical Grasslands-Forrajes Tropicales 7, 267-272.

Torabi, N., Mata, L., Gordon, A., Garrard, G., Wescott, W., Dettmann, P., Bekessy, S.A. (2016) The money or the trees: What drives landholders' participation in biodiverse carbon plantings? Global Ecology and Conservation 7, 1-11.

Tullo, E., Finzi, A., Guarino, M. (2019) Review: Environmental impact of livestock farming and Precision Livestock Farming as a mitigation strategy. Science of The Total Environment 650, 2751-2760.

Tuomisto, H.L., Teixeira de Mattos, M.J. (2011) Environmental Impacts of Cultured Meat Production. Environmental Science & Technology 45, 6117-6123.

Vandermeulen, S., Singh, S., Ramírez-Restrepo, C.A., Kinley, R.D., Gardiner, C.P., Holtum, J.A.M., Hannah, I., Bindelle, J. (2018) *In vitro* assessment of ruminal fermentation, digestibility and methane production of three species of *Desmanthus* for application in northern Australian grazing systems. Crop and Pasture Science 69, 797-807.

Wall, E., Simm, G., Moran, D. (2010) Developing breeding schemes to assist mitigation of greenhouse gas emissions. Animal 4, 366-376.

Weiler, V., Udo, H.M.J., Viets, T., Crane, T.A., De Boer, I.J.M. (2014) Handling multi-functionality of livestock in a life cycle assessment: the case of smallholder dairying in Kenya. Current Opinion in Environmental Sustainability 8, 29-38.

Weinberger, K., Rankine, H., Amanuma, N., Surendra, L., Van Hull, H., Foran, T., Reyes, R., Malik, A., Murray, J.J.U.N., Bangkok: ESCAP (2015) Integrating the three dimensions of sustainable development: A framework and tools.

Westhoek, H., Lesschen, J.P., Rood, T., Wagner, S., De Marco, A., Murphy-Bokern, D., Leip, A., van Grinsven, H., Sutton, M.A., Oenema, O. (2014) Food choices, health and environment: Effects of cutting Europe's meat and dairy intake. Global Environmental Change 26, 196-205.

Wiedemann, S.G., Ledgard, S.F., Henry, B.K., Yan, M.-J., Mao, N., Russell, S.J. (2015) Application of life cycle assessment to sheep production systems: investigating co-production of wool and meat using case studies from major global producers. The International Journal of Life Cycle Assessment 20, 463-476.

Zhang, Y., Gao, Q., Dong, S., Liu, S., Wang, X., Su, X., Li, Y., Tang, L., Wu, X., Zhao, H. (2015a) Effects of grazing and climate warming on plant diversity, productivity and living state in the alpine rangelands and cultivated grasslands of the Qinghai-Tibetan Plateau. The Rangeland Journal 37, 57-65.

Zhang, Y., Huang, D., Badgery, W.B., Kemp, D.R., Chen, W., Wang, X., Liu, N. (2015b) Reduced grazing pressure delivers production and environmental benefits for the typical steppe of north China. Scientific Reports 5, 16434.



gcb_15816_f1.tif







gcb_15816_f3.tif





gcb_15816_f4.tif

LG



gcb_15816_f5.tif





gcb_15816_f6.tif



gcb_15816_f7.tif



gcb_15816_f8.tif



gcb_15816_f9.tif