

# Responses of tundra soil microbial communities to half a decade of experimental warming at two critical depths

Eric R. Johnston<sup>a,b</sup>, Janet K. Hatt<sup>a</sup>, Zhili He<sup>c,d</sup>, Liyou Wu<sup>c,d</sup>, Xue Guo<sup>e</sup>, Yiqi Luo<sup>c,f</sup>, Edward A. G. Schuur<sup>f</sup>, James M. Tiedje<sup>g</sup>, Jizhong Zhou<sup>c,d,e,h</sup>, and Konstantinos T. Konstantinidis<sup>a,i,1</sup>

<sup>a</sup>School of Civil and Environmental Engineering, Georgia Institute of Technology, Atlanta, GA 30332; <sup>b</sup>Biosciences Division, Oak Ridge National Laboratory, Oak Ridge, TN 37830; <sup>c</sup>Institute for Environmental Genomics, University of Oklahoma, Norman, OK 73019; <sup>d</sup>Department of Microbiology and Plant Biology, University of Oklahoma, Norman, OK 73019; <sup>e</sup>State Key Joint Laboratory of Environment Simulation and Pollution Control, School of Environment, Tsinghua University, 100091 Beijing, China; <sup>f</sup>Department of Biological Sciences, Northern Arizona University, Flagstaff, AZ 86011; <sup>g</sup>Center for Microbial Ecology, Michigan State University, East Lansing, MI 48824; <sup>h</sup>Earth Science Division, Lawrence Berkeley National Laboratory, Berkeley, CA 94720; and <sup>i</sup>School of Biological Sciences, Georgia Institute of Technology, Atlanta, GA 30332

Edited by Edward F. DeLong, University of Hawaii at Manoa, Honolulu, HI, and approved June 10, 2019 (received for review January 23, 2019)

Northern-latitude tundra soils harbor substantial carbon (C) stocks that are highly susceptible to microbial degradation with rising global temperatures. Understanding the magnitude and direction (e.g., C release or sequestration) of the microbial responses to warming is necessary to accurately model climate change. In this study, Alaskan tundra soils were subjected to experimental in situ warming by ~1.1 °C above ambient temperature, and the microbial communities were evaluated using metagenomics after 4.5 years, at 2 depths: 15 to 25 cm (active layer at outset of the experiment) and 45 to 55 cm (transition zone at the permafrost/ active layer boundary at the outset of the experiment). In contrast to small or insignificant shifts after 1.5 years of warming, 4.5 years of warming resulted in significant changes to the abundances of functional traits and the corresponding taxa relative to control plots (no warming), and microbial shifts differed qualitatively between the two soil depths. At 15 to 25 cm, increased abundances of carbohydrate utilization genes were observed that correlated with (increased) measured ecosystem carbon respiration. At the 45- to 55-cm layer, increased methanogenesis potential was observed, which corresponded with a 3-fold increase in abundance of a single archaeal clade of the Methanosarcinales order, increased annual thaw duration (45.3 vs. 79.3 days), and increased CH<sub>4</sub> emissions. Collectively, these data demonstrate that the microbial responses to warming in tundra soil are rapid and markedly different between the 2 critical soil layers evaluated, and identify potential biomarkers for the corresponding microbial processes that could be important in modeling.

tundra | metagenomics | permafrost | climate change | soil microbiology

**R**epresenting only  $\sim 16\%$  of Earth's terrestrial surface, northern-latitude permafrost soils and their overlaying active layers harbor an estimated 1,672 to 1,832 Pg of carbon (C), which accounts for ~50% of the global soil organic C (SOC) reservoir (1, 2). This large C stock has accumulated and been preserved for thousands of years, primarily due to low temperatures and frozen conditions, which constrain microbial SOC mineralization (3, 4). Elevated atmospheric greenhouse gas concentrations are increasing global temperatures, and northern-latitude areas are experiencing a rate of warming that is more than twice the global average (5). As a result, regionally widespread and ongoing permafrost thaw is being observed (6-9). It has been estimated that permafrost could recede further by 30 to 70% by the end of the 21st century (10, 11). This alleviation of prior abiotic constraints is expected to stimulate microbial processes resulting in the release of greenhouse gases, primarily CO<sub>2</sub> and CH<sub>4</sub>, which could further exacerbate climate warming (i.e., cause positive feedback) (12-14). However, efforts to delineate microbial processes to improve future climate change predictions are hampered by the enormous complexity and heterogeneity of soil microbial communities and knowledge gaps regarding the microbial metabolic functions and their controls that operate in situ.

Furthermore, performing field warming experiments at northern latitudes is particularly challenging due to the remoteness of these sites and weather conditions. Previous investigations of laboratory-incubated permafrost soils under elevated temperatures reported shifts in community structure and functioning toward increased carbon respiration, even over a short period of a few weeks (15, 16). The relevance of these laboratory findings for in situ processes, however, remains speculative because the laboratory incubations cannot simulate closely the complexity of the natural environment, especially at the deeper soil layers that harbor older carbon stocks. Increasing temperatures and active layer thickness (i.e., an extension of maximum annual thaw depth) also affect other components of the soil, such as redox conditions, water availability, nutrient cycling, and aboveground vegetation

#### Significance

Ongoing permafrost thaw is expected to stimulate microbial release of greenhouse gases, threatening to further exacerbate climate change (cause positive feedback). In this study, a unique field warming experiment was conducted in Interior Alaska to promote surface permafrost degradation while maintaining uniform hydraulic conditions. After 5 winters of experimental warming by ~1 °C, microbial community shifts were observed at the receded permafrost/active layer boundary, which reflected more reduced conditions, including increased methanogenesis. In contrast, increased carbohydrate utilization (respiration) was observed at the surface layer. These shifts were relatable to observed increases in CO<sub>2</sub> and CH<sub>4</sub> release from this study site and the surrounding ecosystem. Collectively, our results demonstrate that microbial responses to warming are rapid and identify potential biomarkers that could be important in modeling.

This article is a PNAS Direct Submission.

Published under the PNAS license.

Data deposition: Metagenome and MAG sequences have been deposited in the European Nucleotide Archive (study ID PRJEB31848). Individual accession IDs for metagenomes and metagenome-assembled genomes can be found in *SI Appendix*, Tables S1 and S5, respectively.

<sup>1</sup>To whom correspondence may be addressed. Email: kostas@ce.gatech.edu.

This article contains supporting information online at www.pnas.org/lookup/suppl/doi:10. 1073/pnas.1901307116/-/DCSupplemental.

Published online July 8, 2019.

Author contributions: E.R.J., Z.H., L.W., Y.L., E.A.G.S., J.M.T., J.Z., and K.T.K. designed research; E.R.J., J.K.H., and X.G. performed research; E.R.J. analyzed data; and E.R.J. and K.T.K. wrote the paper.

The authors declare no conflict of interest.

(17–20). These physicochemical alterations, in turn, can result in additional, indirect effects of warming to soil microbes and their activities. However, such effects remain poorly understood and quantified to date.

More specifically, gradual shifts in vegetative communities can alter soil physical conditions, establish new plant-microbe symbioses, promote oxygen and or methane transit, and change the amount and type of plant material returned to soil (21-24). A combination of soil warming and redox conditions favorable for energy-yielding activities can result in a more rapid release of  $CO_2$  from tundra soils (25, 26). Under lower potential redox conditions, microbes might degrade SOC by gradually using electron acceptors with declining redox potentials, leading to conditions more suitable for methanogenesis. This can result in lower C loss compared with aerobic conditions and yet have greater feedback to climate due to larger CH<sub>4</sub>:CO<sub>2</sub> ratio of emissions (26). Hence, the microbial and environmental features that serve as potential candidates for ecosystem-level evaluation of soil C release are vast and interconnected. Elucidating the key microbial processes and their environmental controls operating in situ requires an integrated approach that takes into account and measures the above parameters. Importantly, which of the microbial processes mentioned above-for example, aerobic/ respiring vs. reducing/methanogenic pathways-prevail in northern latitudes and how fast these processes could be accelerated (or not) by warming remain essentially unknown. An improved understanding of the microbial clades and activities involved could help predict greenhouse gas fluxes and lead to an improved assessment of ecosystem-scale responses to climate warming (27).

To advance these issues, the Carbon in Permafrost Experimental Heating Research (CiPEHR) site was established in a discontinuous permafrost region in Interior Alaska in 2008 and represents the first in situ experiment to promote surface permafrost degradation without delaying spring soil melt (28, 29). In warmed plots, soil is insulated from frigid conditions during the winter months with snow that accumulated on the windward side of plastic mesh snow fences built by our team (SI Appendix, Fig. S1). This results in soil temperatures that are elevated yearround and also in the warming of deep soil, a manipulation not typically achieved with previous tundra warming experiments. Previous reports by our team demonstrated rapid physiological responses of belowground microorganisms after just 2 winters of experimentation at this site. However, communitywide shifts in microbial structure were less discernible at this early stage, perhaps due to a lag between physiological responses and corresponding shifts at the DNA level (30, 31). In this study, shotgun-metagenomic sequencing was used to assess responses of soil community structure after 4.5 y of experimental warming and gain insights into the undergoing in situ processes at 2 critical depths: 15 to 25 cm (preexisting active layer) and 45 to 55 cm (approximately the active layer/permafrost boundary at the initiation of experiment, which has since receded downward). The specific hypotheses tested with these data were: (i) microbial communities at the receded permafrost boundary layer depth will be more sensitive to elevated temperatures; (ii) at both soil depths studied, community functions stimulated by increased temperatures will include traits for the release of  $CO_2$  and/or  $CH_4$ ; and (*iii*) warming-induced changes in the relative abundances of traits involved in  $CO_2$  or  $CH_4$  release will be relatable to increased gas fluxes measured at the CiPEHR site and the surrounding region.

## Results

Environmental Indices. Experimental warming by means of increased winter snow cover resulted in a year-round elevation in soil temperature. For example, warming treatment plots were 1.02 and 1.11 °C warmer than control plots at 15- to 25-cm and 45- to 55-cm soil depths, respectively, during the growing season before sampling (from May to September; sampling took place in May; linear mixed-effects model [LME] used in conjunction with Tukey's honest significant difference [HSD] test, P < 0.05) (Tables 1–3). Experimental warming also increased annual thaw duration from 104.2 to 112.2 d at the 15- to 25-cm depth (7.7% increase; nonsignificant with the LME model, but P < 0.05 using paired t test) and from 45.5 to 79.3 d in the 45- to 55-cm depth (74.4% increase; LME, P < 0.05). The mean thaw depth at the time of sampling in May 2013 was 18.3 and 23.0 cm in control and experimentally warmed plots, respectively (LME, P < 0.05). Experimental warming also increased mean aboveground plant biomass by 25.2%, increased volumetric water content slightly, and increased bulk density at the 45- to 55-cm depth profile (LME, P < 0.05). For aboveground plants, warming decreased lichen biomass by 41.8% on average (paired t test, P < 0.05), and increased the average biomass of Eriophorum vaginatum and Rhododendron tomentosum by 81.2% and 23.3%, respectively (paired t test, P < 0.05).

Broad Microbial Community Indices. On average, 8.1 Gbp (billion base pairs) of sequencing effort per sample was obtained for 12 soil communities representing the 15- to 25-cm soil depth (6 warming, 6 control) and 5.8 Gbp per sample for 12 communities representing the 45- to 55-cm depth (*SI Appendix*, Table S1). Using Nonpareil 3 (options: -T kmer -k 32 -X 100000), the average estimated coverage of the microbial communities assessed with sequencing was 0.52 for 15- to 25-cm and 0.62 for 45- to 55-cm soil communities (SI Appendix, Fig. S2) (32). These coverage values suggest that, beyond the sequencing depth achieved, there is 52% and 62% likelihood that the additional sequences obtained would be redundant with ones already observed. The level of coverage obtained here (52 to 62%) should be appropriate for comparisons (e.g., low false-negative rate), with unsampled diversity presumably representing comparatively rarer taxa (33, 34). Experimental warming resulted in a nearsignificant increase in Nonpareil-derived sequence diversity Nd at the 45- to 55-cm depth, a metric that correlates well with  $\alpha$ -diversity assessed by traditional metrics and is advantageous for shotgun metagenomic data (paired t test, P < 0.1) (SI Ap*pendix*, Fig. S3). MicrobeCensus was used to evaluate whether or not shifts in average genome size of the community due to the experimental warming could account for observed shifts in

Table 1.	Temperature and annual	thaw duration of the ex	perimental groups of the study
----------	------------------------	-------------------------	--------------------------------

Sample group	Annual thaw duration (d)	Growing season 2012 temperate (°C)	Winter 2012/2013 temperature (°C)	May 2013 temperature (°C)
15 to 25 cm Control	$104.2 \pm 1.7^{\circ}$	$2.59 \pm 0.29^{b}$	$-1.15 \pm 0.16^{a}$	$-0.62 \pm 0.1^{a}$
45 to 55 cm Control 45 to 55 cm Warmed	$45.5 \pm 4.6^{a}$ 79.33 ± 5.55 <sup>b</sup>	$1.08 \pm 0.28^{a}$ 2.19 ± 0.39 <sup>b</sup>	$-0.86 \pm 0.07$ $-0.86 \pm 0.13$ <sup>ab</sup> $-0.42 \pm 0.04^{c}$	$-0.20 \pm 0.04$ $-0.73 \pm 0.13^{a}$ $-0.3 \pm 0.05^{b}$

Values are given as the mean  $\pm$  the SEM derived from 6 replicate measurements. Superscript letters are used to distinguish sample groups that were significantly different (adjusted P < 0.05); values with superscript letters differing from letters assigned to other values designate a statistically significant difference between sample groups (i.e., treatment and depth). Statistical significance was determined using a linear mixed-effects model (where experimental fence was treated as a within-subjects factor), in conjunction with Tukey's HSD test.

Johnston et al.

	Table 2.	Summary of soil	physiochemical	measurements for	or the	experimental	groups of the study
--	----------	-----------------	----------------	------------------	--------	--------------	---------------------

Sample group	Soil moisture	Bulk density	Total N (%)	Total C (%)	
15 to 25 cm Control	$0.74\pm0.04^{b}$	$0.25 \pm 0.04^{a}$	1.58 ± 0.11 <sup>b</sup>	$33.4 \pm 3.4^{b}$	
15 to 25 cm Warmed	$0.75 \pm 0.03^{b}$	$0.23 \pm 0.03^{a}$	1.59 ± 0.07 <sup>b</sup>	34.6 ± 1.3 <sup>b</sup>	
45 to 55 cm Control	$0.39 \pm 0.06^{a}$	$0.74 \pm 0.17^{b}$	$0.65 \pm 0.19^{a}$	$16.0 \pm 4.1^{a}$	
45 to 55 cm Warmed	$0.31 \pm 0.03^{a}$	1.19 ± 0.19 <sup>c</sup>	$0.46 \pm 0.11^{a}$	$12.4 \pm 3.0^{a}$	

See Table 1 legend.

 $\alpha$ -diversity or functional gene content (35). Such effects as a result of warming were not observed, but our analysis did reveal smaller genomes, on average, for microbial populations comprising the 45- to 55-cm soil depth compared with those at 15 to 25 cm (average of 4.7 Mbp vs. 6.7 Mbp; LME, P < 0.001) (SI Appendix, Table S1).

Soil Community Taxonomic Composition. The relative 16S rRNA gene-based sequence abundance (metagenome derived) of Ar*chaea* was  $1.69 \pm 0.09\%$  (mean  $\pm$  SEM) and  $2.64 \pm 0.21\%$  in 15to 25-cm and 45- to 55-cm soil depths, respectively (LME, P <0.05) (Fig. 1B and SI Appendix, Table S2). The archaeal community at the 15- to 25-cm depth was mostly represented by phyla Thaumarchaeota, followed by Euryarchaota at 1.11  $\pm$ 0.16% and  $0.42 \pm 0.13\%$ , respectively. For the 45- to 55-cm layer, archaeal 16S rRNA gene sequences were mostly represented by Euryarchaeota and Crenarchaeota at 2.31  $\pm$  0.22% and 0.23  $\pm$ 0.05%, respectively. Warming had no significant effect on the relative abundance of Archaea at the 15- to 25-cm depth (LME, P > 0.1), but increased archaeal abundance by a mean of 39% at 45 to 55 cm (2.21  $\pm$  0.24% in control vs. 3.07  $\pm$  0.27% in warmed soils; LME, P < 0.05). This increase was primarily driven by an increase in the abundance of the archaeal phylum Euryarchaeota  $(1.90 \pm 0.27\% \text{ vs. } 2.73 \pm 0.25\%)$ , and more specifically, the order Methanosarcinales, which was ~3-fold more abundant in experimentally warmed plots at this soil depth  $(0.54 \pm 0.14\% \text{ vs. } 1.61 \pm$ 0.38%; LME, P < 0.05). Notably, a very strong correlation was observed between the relative abundance of Methanosarcinales and annual thaw duration (squared root regression  $R^2 = 0.895$ , P < 0.8950.001) (Fig. 1C). Most of the 18 bacterial phyla with mean relative abundances  $\geq 0.1\%$  across all 24 soil samples differed significantly between 15- to 25-cm and 45- to 55-cm soil communities in relative abundance (excluding only Thermotogae, Lentisphaerae, Armatimonadetes, and Actinobacteria; LME, P < 0.05) (Fig. 1A and SI Appendix, Fig. S4). At 15 to 25 cm, warming only significantly increased the relative abundance of Verrucomicrobia (8.0  $\pm$  0.7% vs. 10.4  $\pm$  0.6% relative abundance in control and warmed samples, respectively; P < 0.05). In contrast, at 45 to 55 cm, experimental warming resulted in significant shifts (P < 0.05) to phyla Proteobacteria  $(23.4 \pm 2.4\% \text{ vs. } 30.4\% \pm 3.4),$ Actinobacteria (31.0  $\pm$  4.1% vs. 20.6  $\pm$  3.0%), and Microgenomates  $(0.2 \pm 0.05\% \text{ vs. } 0.8 \pm 0.2\%)$ .

Consistent with the abovementioned results, at 45 to 55 cm, experimental warming resulted in significant community-wide shifts in taxonomic  $\beta$ -diversity based on abundance weighted Jaccard distances of phylum-level community composition (analysis of similarities [ANOSIM] and multiresponse permutation procedures [MRPP], P < 0.05) (Fig. 2 and Table 4). There was also a significant relationship between  $\beta$ -diversity (derived from either UniFrac or Jaccard distances) and annual thaw duration or depth of thaw at time of sampling (Adonis, P < 0.05). There were no significant relationships between experimental treatment or thaw indices and 16S rRNA gene-based or phylabased evaluations of community structure for samples representing the 15- to 25-cm depth, similar to observations of this depth made after just 2 winters of experimental warming (30).

Shifts in Energy-Yielding Metabolisms with Experimental Warming. In addition to the observed shifts in microbial community phylogenetic structure for 45- to 55-cm metagenomes, experimental warming shifted community functional gene content (summa-rized as Carbohydrate-Active enZYmes [CAZy] families or Kyoto Encyclopedia of Genes and Genomes, or KEGG, Orthology [KO] terms) (ANOSIM, P < 0.05 or P < 0.1) (Fig. 2 and Table 4). Also for this depth,  $\beta$ -diversity based on KO terms was significantly associated with annual thaw duration and measured thaw depth at time of sampling (Adonis, P < 0.05). For 15- to 25-cm soils, warming only significantly altered community  $\beta$ -diversity reflecting the composition of genes involved in carbohydrate metabolism and binding (i.e., CAZy protein families; MRPP, P < 0.05; ANOSIM, P < 0.1), and no significant relationships between functional structure and thaw indices for this depth were identified.

Furthermore, for 15- to 25-cm communities, experimental warming increased the relative proportion of functional genes matching to CAZy reference sequences by 7.9% (LME, P <0.05) (Fig. 3A). This included a 12.0% increase in glycoside hydrolase genes, a 2.6% increase in glucosyltransferase genes, an 8.1% increase in carbohydrate binding genes, an 8.9% increase in carbohydrate esterase gene, and a 20.8% increase in polysaccharide lyase genes (LME, P < 0.05) (Fig. 3 B and C). These results were generally consistent with Gene Ontology (GO) process categories involving the catabolism of various organic matter substrates (Fig. 3D). Of the 25 categories based on these definitions that differed by  $\geq 10\%$  between control and warmed 15- to 25-cm soils, 22 of 25 were more abundant in experimentally warmed metagenomes, with 4 having an adjusted P < 0.05with LME. Similar shifts were not found between warmed and control 15- to 25-cm soils from the 1.5-y collection (SI Appendix, Fig. S5). While warming also shifted community  $\beta$ -diversity of CAZy families at the 45- to 55-cm depth, there was only a significant 3.1% increase in the abundance of glucosyltransferase genes at this depth (LME, P < 0.05).

Warming of the 15- to 25-cm soil layer did not result in any KO term that changed in abundance with an adjusted P < 0.05(using DESeq2) (Datasets S1 and S2). However, of the 3,274 KO terms with abundances adequate for P value assignment in DESeq2, 162 KO terms differed significantly (adjusted P < 0.05) and another 153 differed near-significantly (adjusted P < 0.1) between control and warmed soil community samples at 45 to 55 cm. Also for the 45- to 55-cm depth, the abundances of 490 KO terms increased or decreased significantly across a gradient of annual thaw duration, and another 301 KO terms displayed a near-significant

Table 3. Summary of soil physiochemical measurements for the experimental groups of the study

Sample group	Plant biomass (g)	Thaw depth (cm)	Water table depth (cm)	Volumetric water content (%)
Control plots	$582.3 \pm 24.9^{a}$	$18.3 \pm 0.5^{a}$	$22.96 \pm 2.03^{a}$	$\begin{array}{l} 28.07 \pm 0.14^{a} \\ 29.21 \pm 0.55^{b} \end{array}$
Warmed plots	728.9 $\pm$ 64.6 <sup>b</sup>	23.0 $\pm 1.6^{b}$	$22.29 \pm 4.84^{a}$	

See Table 1 legend.



**Fig. 1.** Taxonomic shifts as an effect of experimental warming. (A) Mean relative abundance of bacterial phyla for each depth  $\times$  treatment combination. Underlying data are based on 165 rRNA gene-encoding fragments recovered from metagenomic datasets. Values represent the abundance of each bacterial phylum as a proportion of the total bacterial community. Only phyla with a mean relative abundance higher than 0.1% across all 24 datasets are displayed. (*B*) Mean relative abundances of archaeal phyla for each depth  $\times$  treatment combination. Underlying data are based on 165 rRNA gene-encoding fragments recovered from metagenomic datasets. Values represent the abundance of all recovered from metagenomic datasets. Values represent the abundance (as a percentage) of each archaeal phylum relative to the total abundance of all recovered bacterial and archaeal 165 rRNA gene fragments (i.e., the total prokaryotic community). Error bars represent the mean  $\pm$  the SEM (n = 6) for the year) for 45- to 55-cm soils. Linear regression was fitted for n = 12 points. Significance of the correlation coefficient was determined using a two-tailed Student's *t*-distribution (r = 0.946, df = 11).

change with thaw duration. These shifts included a broad increase in functions involved in methanogenesis (or potentially, reverse methanogenesis) (36) between control and experimentally warmed communities (Fig. 4A). This evaluation was primarily based on KEGG modules M00567, M00357, M00356, and M00563, corresponding to methanogenesis from CO<sub>2</sub>, acetate, methanol, and methylamines (37) (Fig. 4A). Among these shifts, the largest and consistently significant responses were observed for KO terms specific to methanogenesis from acetate, which increased on average by 170%. There were also strong correlations between the abundance of archaeal order *Methanosarcinales* and KO terms for methanogenesis from acetate ( $R^2 = 0.93$ ; P < 0.01).

Increased temperatures or extended annual thaw of 45- to 55-cm soils also increased the relative abundances of genes involved in dissimilatory sulfate reduction and dissimilatory nitrate reduction to ammonium (adjusted P < 0.05 or 0.1 with thaw or treatment) (Fig. 4B), but resulted in no significant changes in denitrification genes norB and nosZ. The opposite trends were observed for KO terms specific to assimilatory nitrate reduction and assimilatory sulfate reduction (Fig. 4B), implying more reducing redox conditions upon warming and permafrost thawing. An increase in the abundances of genes involved in dissimilatory sulfate reduction was largely attributable to the Proteobacteria order Syntrophobacterales, a clade with many known sulfurreducing taxa, which increased in abundance from 1.0 to 2.5% with experimental warming (P < 0.005). Consistent with this, a recovered metagenome-assembled genome (MAG) belonging to Syntrophobacterales was found to possess genes for respiratory sulfur reduction (dsrAB, aprAB).

DESeq2 analysis revealed statistically significant decreases in the relative abundances of cytochrome *c* oxidase genes with an average 2012 growing season temperature for 45- to 55-cm soils, even when controlling for thaw indices, consistent with more reducing conditions. Multiple linear regression analysis revealed that average summer 2012 temperature and thaw depth had significant independent (controlling for the other variable) as well as additive negative associations with the abundances of cytochrome *c* oxidase genes (multiple  $R^2 > 0.8$ ; P < 0.001) (Fig. 5*A*), and conversely, positive associations with genes involved in methane production from acetate (multiple  $R^2 = 0.85$ ; P < 0.0005) (Fig. 5*B*). The total relative abundance of all CAZy functions in 15- to 25-cm communities, as well as those belonging specifically to carbohydrate binding or glycoside hydrolase modules, had significant positive correlations with cumulative ecosystem respiration in the month following sampling (June 2013) (t-distribution,  $r \ge 0.6$ , P < 0.05) (see also Fig. 5C).

Recovery of MAGs. Assembly and population genome binning led to the recovery of 173 MAGs with a quality score  $\geq 60$  (calculated as completeness –  $5 \times$  contamination based on CheckM) (38, 39) (SI Appendix, Tables S3 and S4). These medium- to high-quality MAGs collectively recruited 17.0% and 24.9% of the short reads, on average, for the 15- to 25-cm and the 45- to 55-cm depth, respectively (using default megablast alignment; ≥97% nucleotide ID and  $\geq$ 100-bp alignment). Dereplication of highly similar MAGs based on FastANI (40) values of  $\geq 95\%$  resulted in the consolidation of 74 nonredundant MAGs. This included 4 nonredundant archaeal MAGs, all derived from 45- to 55-cm soil metagenomes (SI Appendix, Fig. S6). Two archaeal MAGs were identified as belonging to order Methanosarcinales and the other two were assigned to *Methanocellales* based on taxonomic identification with the Microbial Genomes Atlas (MiGA) (SI Appendix, Fig. S7). Each archaeal MAG possessed genes involved in methanogenesis using CO<sub>2</sub>, acetate, and in one case, methanol, as electron acceptors. One Methanosarcinales MAG matched closely to ANME-2d taxa Candidatus Methanoperedens sp. BLZ1 and BLZ2 (87.5% average nucleotide identity [ANI]) (41, 42) and was also assigned to genus Methanoperedens with GTDB-Tk (SI Appendix, Table S4). Similar to the previously recovered Candidatus Methanoperedens taxa, the CiPEHR MAG possessed genes for reverse methanogenesis, nitrate reduction to nitrite (narGH), a cytochrome c nitrite reductase (nrfA), as well as nitrogenase genes nifHDK.

Among bacterial MAGs that shifted in abundance with warming, there was a variable yet large 79% increase in the most dominant bacterial population (making up ~1–2% of the total community) in the 15- to 25-cm active layer with warming (paired *t* test, P < 0.1), which represented a member of the *Acidobacteriaceae* family (53.1% amino acid identity [AAI] with *Candidatus Koribacter versatilis* Ellin345 as determined with MiGA; assigned to family *Koribacteraceae* with GTDB-Tk) (*SI Appendix*, Fig. S8 and Table S4). These results were consistent with smaller, nonsignificant shifts observed after 1.5 y of experimentation for the same taxon. This dominant MAG was previously found to be widespread throughout Alaskan tundra and appears to encode diverse metabolisms for labile and recalcitrant organic matter degradation (31).

July 7, 202'



**Fig. 2.** Functional and phylogenetic shifts as an effect of experimental warming. (*A*) Principle coordinates analysis (PCoA) plot of consolidated KO term annotations. Underlying data are based on abundance-weighted jaccard distance matrix derived from a KO term counts matrix. (*B*) PCoA plot of community phylogenetic composition. Underlying data are a weighted unifrac distance matrix of 16S rRNA gene-encoding fragments recovered with Parallel-META and processed in the QIIME software package, as described in *Materials and Methods*. (C) PCA plot of Swiss-Prot gene annotations. Underlying data are based on a gene count matrix consolidated from Swiss-Prot database references, which underwent variance-stabilizing transformation using the DESeq2 package. The 24 samples shown in the 2D plane of each plot are spanned by their first 2 principal components.

### Discussion

Temperature increases as a result of experimental warming lasted year-round and were moderately uniform between the depth profiles evaluated here (15 to 25 and 45 to 55 cm) (Tables 1 and 2). Climate change and increased temperatures will undoubtedly change numerous important aspects of tundra ecology, such as

thaw depth and plant community factors, which can lead to indirect effects of warming on soil microbial community functioning (18, 25, 43). The magnitude and direction of the microbial responses to these factors could vary considerably between depth profiles. Thus, warming of deep soil extending down to the permafrost boundary layer was a desirable outcome of our experimental warming manipulation. Experimental warming also increased annual thaw duration by 7.7% (+8 d) (relative to control plots) at the 15- to 25-cm soil layer and, in strong contrast, by 74.4% (+33.8 d) at the 45- to 55-cm layer for the year preceding sample collection.

Consistently, greater overall changes in community structure were observed for 45- to 55-cm soil communities. This included an ~3-fold increase in Methanosarcinales, a methanogenic order and the most abundant archaeal clade observed in these soils. The abundance of this Methanosarcinales was highly relatable to annual thaw depth ( $R^2 = \sim 0.9$ , P < 0.001) (Fig. 1C) and accompanied a comprehensive increase in the relative abundances of methanogenesis genes (Fig. 4A). The increase in the relative abundance of these methanogens corresponded to increased CH<sub>4</sub> emissions at the CiPEHR site due to our experimental warming and nearby regions due to ambient warming. For example, CH<sub>4</sub> release was considerably greater from experimentally warmed plots (44) in the same year that soil cores were collected for the current study, and the warming effect from  $CH_4$  emissions is now larger than that from  $CO_2$  in this general area (the Eight Mile Lake region) (45).

Aerobic methane oxidation genes were fairly low in relative abundance, and those that were over the detection limit of our metagenomics effort did not shift in response to warming at either depth (SI Appendix, Fig. S9). It is possible that methane was consumed at a depth not evaluated in this study or by anaerobic methane oxidizing microbiota that remain poorly understood (36, 46). Related to the latter, one of the recovered archaeal MAGs was identified as a close relative to recently described ANME-2d taxa (87.5% ANI to Candidatus Methanoperedens sp. BLZ1) (41), and possessed genes for reverse methanogenesis, dissimilatory nitrate reduction to ammonia (DNRA), and nitrogen fixation (nitrogenase genes nifHDK). Previous reports have demonstrated nitrogen fixation in ANME-2 Archaea (47, 48), but the presence of functions for both DNRA and N<sub>2</sub>-fixation in an ANME-2d genome is unique. This combination of traits (producing ammonium as a waste product of DNRA while also capable of fixation of gaseous N<sub>2</sub> to ammonium, which is energetically expensive) could signify a

Table 4.	The effects of warming a	nd thaw depth or du	ration on soil microbial comr	munity functional an	d phylogenetic structure
----------	--------------------------	---------------------	-------------------------------	----------------------	--------------------------

		Adonis		ANOSIM	MRPP	
Underlying data	Distance metric	Thaw time	Thaw depth	Warming treatment	Warming treatment	
15- to 25-cm Soil depth						
16S-Based OTUs	Weighted Unifrac	0.697	0.488	0.587	0.187	
Phyla composition	Bray–Curtis	0.463	0.165	0.398	0.442	
	Abundance-weighted Jaccard	0.455	0.479	0.915	0.747	
KO terms	Bray–Curtis	0.416	0.761	0.894	0.769	
	Abundance-weighted Jaccard	0.491	0.576	0.440	0.119	
CAZy families	Bray–Curtis	0.232	0.441	0.543	0.346	
	Abundance-weighted Jaccard	0.330	0.284	0.083	0.047	
45- to 55-cm Soil depth						
16S-Based OTUs	Weighted Unifrac	0.011	0.006	0.186	0.120	
Phyla composition	Bray–Curtis	0.018	0.002	0.131	0.087	
	Abundance-weighted Jaccard	0.045	0.053	0.043	0.039	
KO terms	Bray–Curtis	0.048	0.069	0.091	0.139	
	Abundance-weighted Jaccard	0.010	0.006	0.099	0.068	
CAZy families	Bray–Curtis	0.170	0.152	0.014	0.142	
	Abundance-weighted Jaccard	0.523	0.445	0.095	0.438	

Permutation procedure (MRPP) and analysis of similarity (ANOVA) were used to test the effect of experimental warming, and permutational multivariate ANOVA (Adonis) was used to test the effect of thaw depth (at time of sampling; in centimeters) and the annual duration of thaw (in days of the year) on each community metric. Numerical values represent probability scores (i.e., P values) resulting from each test. P < 0.1 are bolded to highlight significant and near-significant results.



**Fig. 3.** Shifts in carbohydrate metabolism genes as an effect of experimental warming. Relative abundances of (A) all CAZy annotations, and CAZy modules (*B*) glycoside hydrolases (GH), glycosyl transferases (GT), and (*C*) auxiliary activities (AA), carbohydrate binding (CBM), carbohydrate esterases (CE), and polysaccharide lyases (PL). Underlying values represent the mean relative abundance of each category for each sample group (i.e., treatment × depth). The abundances of CAZy functions were determined by normalizing the number of annotations matching to each broad definition by the number of annotations matching to the more comprehensive Swiss-Prot database. Letters distinguish sample groups that were significantly different (adjusted *P* < 0.05); that is, values with letters differing from letters assigned to other groups designate a statistically significant difference between groups. Significance was determined using a LME model where experimental fence was treated as a within-subjects factor, in conjunction with Tukey's HSD test. (*D*) Shifts in the relative abundances of catabolic pathways were determined by matching Swiss-Prot references to GO Biological Process terms, and dividing the number of annotations. Bars reflect the mean relative abundance for each sample group (unwarmed, warmed) at 15 to 25 cm (*n* = 6) and error bars represent the SEM. Only pathways that differed between warmed and unwarmed soils by  $\geq$ 10%, on average, are shown.

high degree of metabolic versatility or alternative functions associated with these genes. Related to this, previous research has provided some evidence of ANME taxa capable of both forward and reverse methanogenesis (49); hence, it remains unclear what role the ANME-2d population has in methane fluxes in these tundra soils.

For the last decade (2008-2017), the majority of tundra habitats have exhibited a 2 to 4.2 °C temperature increase above the 1950–1980 base period during cold seasons of the Northern Hemisphere (November to April) (50, 51). Despite an increase in background temperature that has continued to recede the permafrost boundary layer in warmed and unwarmed plots, a rapid and clear response of methanogenic taxa was nonetheless observed in experimentally warmed plots. Elevated temperatures and increased thaw were associated with independent as well as additive declines in the abundances of O<sub>2</sub>-specific respiratory metabolisms, such as cytochrome c oxidase genes (Fig. 5A). The opposite trend was found for genes involved in methane production from acetate (Fig. 5B), reflecting a shift toward lower reduction potential. It is possible that accelerated warming caused by our experimental set up caused a more thorough depletion of electron acceptors compared with gradual warming (i.e., recent ambient), due to a smaller period over, which the acceptors can be replenished, resulting in a greater CH<sub>4</sub>:CO<sub>2</sub> ratio of emissions. A decline in redox conditions was also evidenced by greater relative abundances of DNRA and dissimilatory sulfate reduction genes (Fig. 4B), and associated taxa, such as Proteobacteria order Syntrophobacterales. Accompanying declines in assimilatory nitrogen and sulfur reduction genes observed under experimentally warmed conditions at 45 to 55 cm could be due to enhanced  $NH_4^+$  and  $S^{2-}$  production from dissimilatory reduction pathways, and thus a selection of traits for direct acquisition of these reduced compounds (Fig. 4B).

Warming also increased the biomass of *E. vaginatum* by 81%, a plant species that excretes organic acids such as acetate that can fuel CH<sub>4</sub> production, transit, and release (23). Another plant

species known to fuel methanogens, *Carex rostrata* (23), is common in this field location, albeit with lower biomass. It is possible that a greater response observed for genes specific to methanogenesis from acetate compared with other methanogenesis pathways (Fig. 4*A*) is attributable at least in part to these types of plant–microbe interactions. However, the associations between plant and methanogens resulting in greater CH<sub>4</sub> emission from tundra habitats remains elusive (52, 53), and thus are topics worthy of consideration in future studies.

At 15 to 25 cm, warming increased the total proportion of community functional genes involved in carbohydrate metabolism (Fig. 3A), including glycoside hydrolases, glycosyltransferases, carbohydrate esterases, carbohydrate binding, and polysaccharide lyases (Fig. 3 B and C). Furthermore, the relative abundances of these carbohydrate metabolism categories were relatable to increased ecosystem respiration as a result of experimental warming in the month following sampling (June 2013) (Fig. 5C)  $(r \ge 0.6, P < 0.05)$ . An early assessment of this depth profile after 1.5 y of experimental warming found a similar response using GeoChip that was undetected or less obvious with metagenomic analysis (30). Thus, the response observed here is consistent with earlier observations, but also implies an ongoing change over time (SI Appendix, Fig. S5). Consolidation of CAZy reference sequences into family definitions also revealed a significant relationship between warming treatment and the relative composition (i.e.,  $\beta$ -diversity) of these functions (Table 4). Moderately strong associations between the fraction of genes involved in carbohydrate metabolism and the average 15- to 25-cm soil temperature during the same month for which soils were collected (May) ( $R^2 = 0.521$ ; P < 0.01) and the average temperature of the preceding winter season ( $R^2 = 0.478$ ; P < 0.05) were also observed. These direct associations between the community fraction of genes involved in carbohydrate metabolism and soil temperature were much stronger than correlations from other broad environmental measure displayed in Tables 1-3, including plant biomass ( $R^2 = 0.179$  or less; P > 0.1), which increased by

Johnston et al.



**Fig. 4.** Shifts in microbial energy-generating pathways as an effect of experimental warming. (*A*) Heatmap showing the relative abundances of genes involved in methanogenesis (or potentially, reverse methanogenesis) for 45- to 55-cm depth soil metagenomes. KO terms under the "Core methanogenesis genes" category represent those used by all methanogens. Other categories and KO terms refer to pathways involved in the usage of different electron acceptors to sustain methanogenesis, which are variable between methanogenic taxa. To emphasize differences between datasets, the values for each gene were normalized by the mean relative abundance of all 12 of the 45- to 55-cm soil metagenomes (by rows). Subplots on the right of the heatmap represent the  $log_2$  fold-difference calculated for each gene, and are colored by whether differences between control and experimentally warmed soils were statitistical significant (*Right*) or if there was a significant association between gene relative abundance and annual thaw duration (*Left*) (dark gray = adjusted P < 0.05; light gray = nonadjusted P < 0.05; white = nonsignificant). Subplot on the top of the heatmap represents the annual thaw duration (i.e., days of the year soil was thawed) for each plot represented by the corresponding metagenomes. (*B*) Heatmaps showing the relative gene abundances for assimilatory and dissimilatory nitrate and sulfate reduction. Underlying data for *A* and *B* are communities respresenting 45- to 55-cm soil samples.

25.2% in experimentally warmed plots (Table 3) (paired *t* test P < 0.05). These results of stimulated functions involved in organic C turnover as a direct response to warming are somewhat surprising, as recent C releases in an adjacent area were found to result mostly from new SOC sources (17). This could suggest that even with enhanced vegetative production, increases in the relative abundances of genes involved in carbohydrate metabolism are nonetheless more attributable to soil temperature, at least in frigid locations where microbial catabolic functions are constrained by cold conditions.

#### **Conclusions and Future Perspectives**

Our evaluation revealed a much greater susceptibility of soil communities at the recently receded permafrost/active layer boundary to further thaw and temperature increase; these responses contrasted with warming-induced changes to shallower, preexisting active layer communities. Community metabolic shifts in the deeper layer strongly indicated a stimulation of methanogens, particularly those using acetate for  $CH_4$  production, which was consistent with independent measurements

of methane emissions from the same site (44). These effects were not solely attributable to increased thaw depth or duration, but were in conjunction with increased temperatures. If accelerated warming results in more thorough depletion of electron acceptors than would occur under a more gradual thaw, it could result in greater CH<sub>4</sub>:CO<sub>2</sub> of emissions, and thus stronger positive feedback to climate change. Meanwhile, increasing  $CH_4$  emissions observed in this area now outweigh the warming potential of CO<sub>2</sub> emissions and offsets C uptake by local plants (44, 45). Furthermore, northern-latitude areas have experienced a rate of warming over the past several decades that is much greater than the global average. If responses under rapid vs. gradual warming are dissimilar in potential CH4 release, this could imply that inferences made from natural thaw gradients (54) or geological records may not serve as adequate predictors of future CH<sub>4</sub> release from tundra soils if warming continues at a rapid pace. Shifts observed at the 15- to 25-cm layer were distinctly attributable to functions involved in carbohydrate metabolisms, were most attributable to the direct effect of elevated temperatures, and were also relatable to enhanced ecosystem respiration.



Fig. 5. Shifts in energy generating metabolisms with temperature and thaw depth in 45- to 55-cm soil communities. (A) Independent and additive effects of thaw depth and average summer temperature on the relative abundances of cytochrome c oxidase genes and (B) genes involved in methane generation from acetate. (C) Cumulative June 2013 ecosystem respiration for each experimental plot. Three-dimensional plots were made using the R package "Scatterplot3d" and illustrate relationships between average summer soil temperatures, thaw depth at time of sampling, and the abundance of energy generating metabolisms summarized as KO terms (the title of each plot reflects the KO identifier, as well as the common gene identifier). R function Im() was used to fit multiple variables to determine assess their independent and combinatorial effects. Individual data points are colored by average summer 2012 temperature, which proceed from blue to red to reflect low vs. comparatively higher soil temperatures.

While an increase in the abundance of methanogenic taxa and genes certainly implies that these CH4-generating taxa were active at some point, numerous environmental conditions and biological factors (i.e., plants, methanotrophs) ultimately determine methane emission vs. consumption. One archaeal MAG closely matching the recently described ANME-2d taxa possessed genes for reverse methanogenesis, DNRA, and nitrogen fixation. This combination of traits highlights uncertainty regarding its role in N and C flux in tundra systems. Hence, there remains a need to understand how susceptible these soils are to CH<sub>4</sub> loss and the microbial mechanisms involved. Assessments of in situ microbial activity, such as through the use of proteomics and transcriptomics of soils (16, 55, 56), can offer such insights regarding whole-community or individual-taxon functioning. The emerging breadth of approaches enabling a more comprehensive investigation of soil microbiota should further serve to unravel the complex responses and interactions between ecological attributes in warming tundra ecosystems. An improved understanding of warming tundra ecosystems will greatly improve our understanding of their contributions to ongoing climate change. The gene and genome sequences reported here should help facilitate proteomics and primer design for PCR assays that can be used to precisely monitor the dynamics of responsive populations in tundra and elsewhere, and further corroborate the results reported herein.

## **Materials and Methods**

Study Site Description and Sample Collection. The CiPEHR site was established at a moist acidic tundra area in September 2008 in Interior Alaska near Denali National Park in the Eight Mile Lake region (63°52′59″N, 149°13′32″W) in a discontinuous permafrost region where permafrost thaw has been observed in the past several decades. Experimental design and site description have been described in detail previously (28); see also *SI Appendix*, Fig. S1 for a diagram and photos of the experimental fences. In brief, 3 experimental blocks were located ~100 m away from each other. In each block, 2 snow fences were constructed about 5-m apart in the winter. Snow fences were marked out of plastic mesh netting with ~2-inch<sup>2</sup> holes. The winter warming treatment plots were located 5-m back from the leeward side of the snow fences, while plots at the windward side of snow fences served as paired

EARTH, ATMOSPHERIC, ND PLANETARY SCIENCES

hydraulic conditions between winter warming and control treatments. Air temperature from 2004 to 2013 ranged from a monthly average of  $-18.0 \pm 1.8$  °C in January to  $13.4 \pm 0.5$  °C in July, with an average annual temperature of  $-2.7 \pm 0.4$  °C. The mean annual precipitation was 378 mm and the mean growing season precipitation from 2004 to 2013 was 216  $\pm$ 24 mm (44). Only C<sub>3</sub> plant species were observed in this area; dominant species include E. vaginatum, Vaccinium uliginosum, some other vascular species, nonvascular feather moss, and lichen. In the experimental plots, the upper 45 to 65 cm of soil was rich in organic C materials and below was mineral soil with a mixture of glacial till and windblown loess. The active layer depth was about 50 cm at initiation of the experiment but has since expanded to lower depths (44, 45). Twelve soil cores, 6 each from warming treatment and control plots, were collected using electric drills in the beginning of the 2010 and 2013 growing season (May), 2 (1.5 y) and 5 winters (4.5 y) after the start of the winter warming manipulation. Cumulative ecosystem respiration for the month following sampling was based on halfhourly chamber measurements used to evaluate net ecosystem exchange, where nighttime net ecosystem exchange was taken as ecosystem respiration (57). Additional information on site monitoring and characterization of environmental indices can be found in SI Appendix.

snow was removed in the spring before snow melt to maintain uniform

Soil DNA Sequencing and Read Processing. Soil DNA extraction, metagenome library preparation, and sequencing of 4.5-y soil samples were performed similarly to methods used for soils collected after 1.5 y (31); see *SI Appendix* for details. The 4.5-y metagenome datasets were deposited in the European Nucleotide Archive under study no. PRJEB31848 (58). Accession IDs corresponding to each sample metagenome are provided in *SI Appendix*, Table S1. Metagenomes representing the 1.5-y sample collection are available under ENA project ID PRJEB10725 (31). Metagenomic paired-end reads were merged using PEAR (59) (options: -p 0.001). All merged and nonmerged reads were then quality-trimmed with the SolexaQA package (60) (options: -h 17;  $\geq$ 98% accuracy per nucleotide position). Trimmed sequences used downstream for functional annotation or taxonomic assignment were truncated to 150 bp to avoid read-length biases.

Soil Community 16S rRNA Gene Analysis. For the assessment of taxonomic composition, 16S and 23S rRNA gene fragments were first recovered from metagenomes using SortMeRNA (61). The relative proportion of sequences matching to bacterial vs. archaeal rRNA gene sequences from SortMeRNA

was used to estimate the relative abundances of either domain. The relative abundances of bacterial phyla were determined with Parallel-META 2.0 (62), using SortMeRNA output sequences as input. Archaeal phyla abundances were determined by aligning SortMeRNA output sequences against the SILVA SSU database (v132) (63) using BLASTN alignment (BLAST+ v2.2.28; options: -word\_size 18) (64). Matches to archaeal sequences with  $\geq$ 80% nucleotide identity and  $\geq$ 100-bp alignment length were retained. To assess community phylogenetic structure, SortMeRNA output sequences were used with the QIIME 1 software package (65) for closed-reference OTU picking with UCLUST (options: -m uclust\_ref -C -z) (66) against the SILVA SSU database dereplicated at 99% nucleotide identity. Aligned SILVA reference sequences represented in the OTU table were then used to construct a phylogenetic tree using FastTree (67). The resulting phylogenetic tree and OTU count table was then used to generate a weighed UniFrac distance matrix summarizing pairwise phylogenetic distances between communities.

Soil Community Functional Gene Assessment. Short nucleotide sequences were searched against the Swiss-Prot (downloaded on November 27, -2016) (68) and CAZy (downloaded on July 15, 2016) (69) reference databases using DIAMOND BLASTX alignment (options: -k 1 -e 1E-5-sensitive) (70). Matches to Swiss-Prot or CAZy reference sequences with a bit score ≥55 were retained for further analysis. Independent count matrices were made to summarize the results of alignment against either database. A reference data file associating KO functions (KO terms) (37) to Swiss-Prot database entries obtained from https://www.uniprot.org/downloads was used to convert Swiss-Prot annotation counts into a more consolidated count matrix of KO terms (available as Dataset S1). Swiss-Prot annotations were also consolidated into GO categories as described previously (31). Annotations to CAZy reference sequences were consolidated into a count matrix of structurally related catalytic and carbohydrate-binding families. They were also consolidated further into the 6 broad functional modules covered in the CAZy database, including glycoside hydrolases, glycosyl transferases, auxiliary activities, carbohydrate binding, carbohydrate esterases, and polysaccharide lyases. For each sample metagenome, the abundances of summarized CAZy definitions, as proportions of all functional genes, were determined by normalizing the number of annotations matching to each broad definition by the number of annotations matching to the more functionally comprehensive Swiss-Prot database as well as the total number of sequences originally used as query for functional annotation. For the summarized count matrices of KO terms or CAZy enzyme families, compositional dissimilarity between samples was calculated with Bray-Curtis and abundance-weighted Jaccard distance metrics.

**DNA Assembly, Genome Binning, and MAG Quality Assessment.** Metagenomic sequences were assembled with IDBA-UD (71) (options:-mink 43-maxk 123-step 4-min\_contig 300). Resulting contigs ≥ 2.5 kbp were used with MetaBAT2 (options:-minCVSum 10) (72) and MaxBin2 (options:-min\_contig\_length 2000) (73) to recover microbial population genomes. Before binning, Bowtie 2 was used to align short-read sequences to embled contigs (options:-very-fast) (74) and SAMtools was used to sort and convert SAM files to BAM format (75). Sorted BAM files were then used to calculate the coverage (mean representation) of each contig in each sample metagenome. The quality of each resulting MAG was evaluated with CheckM v1.0.3 taxonomy workflow for *Bacteria* and *Archaea*, separately (39). The result from either evaluation (i.e., taxonomy workflow for *Chronea or Bacteria*) with the highest estimated completeness was retained for each MAG. MAGs with a quality score ≥60 were retained (from ref. 38; calculated as the estimated complete-

- C. Tarnocai et al., Soil organic carbon pools in the northern circumpolar permafrost region. Glob. Biogeochem. Cycles 23, GB2023 (2009).
- C. Mu et al., Editorial: Organic carbon pools in permafrost regions on the Qinghai– Xizang (Tibetan) Plateau. Cryosphere 9, 479–486 (2015).
- E. A. G. Schuur et al., Vulnerability of permafrost carbon to climate change: Implications for the global carbon cycle. *Bioscience* 58, 701–714 (2008).
- C. E. Hicks Pries, E. A. G. Schuur, K. G. Crummer, Holocene carbon stocks and carbon accumulation rates altered in soils undergoing permafrost thaw. *Ecosystems* 15, 162– 173 (2012).
- IPCC, "Near-term climate change: projections and predictability" in Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change, T. F. Stocker et al., Eds. (Cambridge University Press, Cambridge, UK, 2013), pp. 953–1028.
- M. T. Jorgenson, C. H. Racine, J. C. Walters, T. E. Osterkamp, Permafrost degradation and ecological changes associated with a warming climate in central Alaska. *Clim. Change* 48, 551–579 (2001).
- D. M. Lawrence, A. G. Slater, A projection of severe near-surface permafrost degradation during the 21st century. *Geophys. Res. Lett.* 32, L24401 (2005).

ness –  $5 \times$  contamination). FastANI (40) was used to estimate the ANI between MaxBin 2 and MetaBAT 2 generated MAGs. For redundant MAGs obtained independently from either binning method that matched at >95% ANI and were derived from the same assembled metagenome, the MAG version with the highest quality score or greatest overall size was retained while the other was discarded. Protein-encoding genes from MAG contig sequences were predicted with Prodigal (76) and the resulting nucleotide sequences were searched against the Swiss-Prot database reference sequences using DIAMOND BLASTX alignment as described above. Select MAGs were also processed using the MiGA webserver for taxonomic assignment (77). MAGs uploaded to the MiGA online webserver are freely available under MAGs project "AK Tundra" for browsing of their amino acid identity relatedness, taxonomic classification against named taxa, and quality of genome sequence (http://enve-omics. ce.gatech.edu:3000/projects/22). MAG taxonomic assignments with MiGA and European Nucleotide Archive accession IDs are provided in SI Appendix, Table S5. GTDB-Tk was also used for taxonomic assignment of MAGs (SI Appendix, Table S4) and for generating phylogenetic trees for bacterial (Dataset S3) and archaeal (Dataset S4) MAGs (78).

Statistical Analyses. With the R package nlme (79), a LME model treating experimental fence as a within-subjects factor was used to evaluate differences between experimental groups (15- to 25-cm unwarmed, 15- to 25-cm warmed, 45- to 55-cm unwarmed, 45- to 55-cm warmed) based on their environmental indices, the relative abundances of Bacteria, Archaea, and their phyla, and the relative proportions of summarized CAZy modules or broad GO biological process terms. Tukey's HSD test was used as a post hoc analysis for assessing statistical significance of differences in the means of 4 treatments (15- to 25-cm control, 15- to 25-cm warming, 45- to 55-cm control, and 45- to 55-cm warming). ANOSIM and MRPP were used to evaluate the significance of compositional differences between warmed and control community functional and phylogenetic structure at either depth. Adonis was used to identify statistically significant associations between sample distances and pertinent environmental indices (e.g., thaw depth at sampling, annual thaw duration). Significance of correlations between metadata and certain community measures (metagenome annotations) was assessed using a 2-tailed t-distribution. Tables with raw counts of KO terms were processed with the DESeg2 software package (80) to identify significant, differentially abundant functions between treatments within a depth (where fence was treated as a within-subjects factor using design = ~ Fence + Treatment) or using annual thaw duration as a continuous variable. Benjamini-Hochberg P value correction (81) was used to account for falsediscovery rate arising from multiple comparisons. DESeg2 results are provided as Dataset S2. The R package Superheat (82) was used to illustrate the relative abundances of annotations in conjunction with metadata and statistical results. The lm() function in R was used to perform multiple linear regression to evaluate potential interactive effects between thaw depth and average summer 2012 temperature on the relative abundances of genes involved in energy-yielding metabolisms. Paired t test was used to compare certain ecological factors that did not involve a depth component.

ACKNOWLEDGMENTS. This research was supported by US Department of Energy Award DE-SC0004601 (to J.Z., Z.H., L.W., Y.L., E.A.G.S., J.M.T., and K.T.K.) and US National Science Foundation Awards 1356288 and 1759831 (to K.T.K.). We thank Joel Kostka for his feedback on this manuscript as well as the Partnership for an Advanced Computing Environment at the Georgia Institute of Technology, which enabled the computational tasks associated with this study.

- T. E. Osterkamp et al., Physical and ecological changes associated with warming permafrost and thermokarst in Interior Alaska. *Permafr. Periglac. Process.* 20, 235–256 (2009).
- V. E. Romanovsky, S. L. Smith, H. H. Christiansen, Permafrost thermal state in the polar Northern Hemisphere during the international polar year 2007-2009: A synthesis. *Permafr. Periglac. Process.* 21, 106–116 (2010).
- E. A. G. Schuur, B. Abbott, Climate change: High risk of permafrost thaw. Nature 480, 32–33 (2011).
- D. M. Lawrence, A. G. Slater, S. C. Swenson, Simulation of present-day and future permafrost and seasonally frozen ground conditions in CCSM4. J. Clim. 25, 2207–2225 (2012).
- M. Heimann, M. Reichstein, Terrestrial ecosystem carbon dynamics and climate feedbacks. *Nature* 451, 289–292 (2008).
- 13. E. A. G. Schuur *et al.*, Expert assessment of vulnerability of permafrost carbon to climate change. *Clim. Change* **119**, 359–374 (2013).
- B. W. Abbott *et al.*, Biomass offsets little or none of permafrost carbon release from soils, streams, and wildfire: An expert assessment. *Environ. Res. Lett.* **11**, 034014 (2016).

- R. Mackelprang et al., Metagenomic analysis of a permafrost microbial community reveals a rapid response to thaw. Nature 480, 368–371 (2011).
- M. J. L. Coolen, W. D. Orsi, The transcriptional response of microbial communities in thawing Alaskan permafrost soils. Front. Microbiol. 6, 197 (2015).
- E. A. G. Schuur *et al.*, The effect of permafrost thaw on old carbon release and net carbon exchange from tundra. *Nature* **459**, 556–559 (2009).
- L. D. Hinzman et al., Trajectory of the Arctic as an integrated system. Ecol. Appl. 23, 1837–1868 (2013).
- D. A. Lipson et al., Metagenomic insights into anaerobic metabolism along an Arctic peat soil profile. PLoS One 8, e64659 (2013).
- F. Keuper et al., A frozen feast: Thawing permafrost increases plant-available nitrogen in subarctic peatlands. Glob. Change Biol. 18, 1998–2007 (2012).
- H. P. Bais, T. L. Weir, L. G. Perry, S. Gilroy, J. M. Vivanco, The role of root exudates in rhizosphere interactions with plants and other organisms. *Annu. Rev. Plant Biol.* 57, 233–266 (2006).
- P. Grogan, S. Jonasson, Controls on annual nitrogen cycling in the understory of a subarctic birch forest. *Ecology* 84, 202–218 (2003).
- L. Ström, M. Mastepanov, T. R. Christensen, Species-specific effects of vascular plants on carbon turnover and methane emissions from wetlands. *Biogeochemistry* 75, 65– 82 (2005).
- 24. C. E. Hicks Pries, E. A. G. Schuur, K. G. Crummer, Thawing permafrost increases old soil and autotrophic respiration in tundra: Partitioning ecosystem respiration using  $\delta$ <sup>(13)</sup> C and  $\Delta$ <sup>(14)</sup> C. *Glob. Change Biol.* **19**, 649–661 (2013).
- B. Elberling et al., Long-term CO2 production following permafrost thaw. Nat. Clim. Chang. 3, 890–894 (2013).
- H. Lee, E. A. G. Schuur, K. S. Inglett, M. Lavoie, J. P. Chanton, The rate of permafrost carbon release under aerobic and anaerobic conditions and its potential effects on climate. *Glob. Change Biol.* 18, 515–527 (2012).
- C. K. McCalley et al., Methane dynamics regulated by microbial community response to permafrost thaw. Nature 514, 478–481 (2014).
- S. M. Natali et al., Effects of experimental warming of air, soil and permafrost on carbon balance in Alaskan tundra. Glob. Change Biol. 17, 1394–1407 (2011).
- S. M. Natali, E. A. G. Schuur, R. L. Rubin, Increased plant productivity in Alaskan tundra as a result of experimental warming of soil and permafrost: Increased plant productivity in Alaskan tundra. J. Ecol. 100, 488–498 (2012).
- K. Xue et al., Tundra soil carbon is vulnerable to rapid microbial decomposition under climate warming. Nat. Clim. Chang. 6, 595–600 (2016).
- E. R. Johnston *et al.*, Metagenomics reveals pervasive bacterial populations and reduced community diversity across the Alaska tundra ecosystem. *Front. Microbiol.* 7, 579 (2016).
- L. M. Rodriguez-R, S. Gunturu, J. M. Tiedje, J. R. Cole, K. T. Konstantinidis, Nonpareil 3: Fast estimation of metagenomic coverage and sequence diversity. *mSystems* 3, e00039-18 (2018).
- L. M. Rodriguez-R, K. T. Konstantinidis, Estimating coverage in metagenomic data sets and why it matters. ISME J. 8, 2349–2351 (2014).
- X. Zhang, E. R. Johnston, L. Li, K. T. Konstantinidis, X. Han, Experimental warming reveals positive feedbacks to climate change in the Eurasian Steppe. *ISME J.* 11, 885– 895 (2017).
- S. Nayfach, K. S. Pollard, Average genome size estimation improves comparative metagenomics and sheds light on the functional ecology of the human microbiome. *Genome Biol.* 16, 51 (2015).
- S. J. Hallam et al., Reverse methanogenesis: Testing the hypothesis with environmental genomics. Science 305, 1457–1462 (2004).
- M. Kanehisa, Y. Sato, M. Kawashima, M. Furumichi, M. Tanabe, KEGG as a reference resource for gene and protein annotation. *Nucleic Acids Res.* 44, D457–D462 (2016).
- D. H. Parks et al., Recovery of nearly 8,000 metagenome-assembled genomes substantially expands the tree of life. Nat. Microbiol. 2, 1533–1542 (2017).
- D. H. Parks, M. Imelfort, C. T. Skennerton, P. Hugenholtz, G. W. Tyson, CheckM: Assessing the quality of microbial genomes recovered from isolates, single cells, and metagenomes. *Genome Res.* 25, 1043–1055 (2015).
- C. Jain, L. M. Rodriguez-R, A. M. Phillippy, K. T. Konstantinidis, S. Aluru, High throughput ANI analysis of 90K prokaryotic genomes reveals clear species boundaries. *Nat. Commun.* 9, 5114 (2018).
- A. Arshad et al., A metagenomics-based metabolic model of nitrate-dependent anaerobic oxidation of methane by Methanoperedens-like Archaea. Front. Microbiol. 6, 1423 (2015).
- S. Berger, J. Frank, P. Dalcin Martins, M. S. M. Jetten, C. U. Welte, High-quality draft genome sequence of "Candidatus Methanoperedens sp." strain BLZ2, a nitratereducing anaerobic methane-oxidizing archaeon enriched in an anoxic bioreactor. Genome Announc. 5, e01159-17 (2017).
- S. A. Sistla, J. P. Schimel, Seasonal patterns of microbial extracellular enzyme activities in an arctic tundra soil: Identifying direct and indirect effects of long-term summer warming. Soil Biol. Biochem. 66, 119–129 (2013).
- S. M. Natali et al., Permafrost thaw and soil moisture driving CO 2 and CH 4 release from upland tundra. J. Geophys. Res. Biogeosci. 120, 525–537 (2015).
- M. A. Taylor, G. Celis, J. D. Ledman, R. Bracho, E. A. G. Schuur, Methane efflux measured by eddy covariance in Alaskan upland tundra undergoing permafrost degradation. J. Geophys. Res. Biogeosci. 123, 2695–2710 (2018).
- R. K. Thauer, S. Shima, Methane as fuel for anaerobic microorganisms. Ann. N. Y. Acad. Sci. 1125, 158–170 (2008).
- A. E. Dekas, S. A. Connon, G. L. Chadwick, E. Trembath-Reichert, V. J. Orphan, Activity and interactions of methane seep microorganisms assessed by parallel transcription and FISH-NanoSIMS analyses. *ISME J.* 10, 678–692 (2016).

- V. Krukenberg et al., Gene expression and ultrastructure of meso- and thermophilic methanotrophic consortia. Environ. Microbiol. 20, 1651–1666 (2018).
- F. Beulig, H. Røy, S. E. McGlynn, B. B. Jørgensen, Cryptic CH<sub>4</sub> cycling in the sulfatemethane transition of marine sediments apparently mediated by ANME-1 archaea. *ISME J.* **13**, 250–262 (2019).
- J. Hansen, R. Ruedy, M. Sato, K. Lo, Global surface temperature change. *Rev. Geophys.* 48, RG4004 (2010).
- GISTEMP Team (2018) GISS Surface Temperature Analysis (GISTEMP). NASA Goddard Institute for Space Studies. https://data.giss.nasa.gov/gistemp. Dataset accessed 11 July 2018.
- Y. Wang et al., Influence of plant species and wastewater strength on constructed wetland methane emissions and associated microbial populations. *Ecol. Eng.* 32, 22– 29 (2008).
- 53. J. Kao-Kniffin, D. S. Freyre, T. C. Balser, Methane dynamics across wetland plant species. Aquat. Bot. 93, 107–113 (2010).
- B. J. Woodcroft et al., Genome-centric view of carbon processing in thawing permafrost. Nature 560, 49–54 (2018).
- J. Hultman et al., Multi-omics of permafrost, active layer and thermokarst bog soil microbiomes. Nature 521, 208–212 (2015).
- E. R. Johnston *et al.*, Phosphate addition increases tropical forest soil respiration primarily by deconstraining microbial population growth. *Soil Biol. Biochem.* **130**, 43– 54 (2019).
- M. Mauritz et al., Eight Mile Lake Research Watershed, Carbon in Permafrost Experimental Heating Research (CiPEHR): Half-hourly growing season, chamber-based, CO2 flux data, 2009-2017. https://portal.edirepository.org/nis/metadataviewer?packageid=knb-lter-bnz.481.19. Accessed 15 June 2018.
- E. R. Johnston, Responses of tundra soil microbial communities to half a decade of experimental in-situ warming at two critical depths (CiPEHR Alaska site). European Nucleotide Archive. https://www.ebi.ac.uk/ena/data/view/PRJEB31848. Deposited 7 May 2019.
- 59. J. Zhang, K. Kobert, T. Flouri, A. Stamatakis, PEAR: A fast and accurate illumina paired-end reAd mergeR. *Bioinformatics* **30**, 614–620 (2014).
- M. P. Cox, D. A. Peterson, P. J. Biggs, SolexaQA: At-a-glance quality assessment of Illumina second-generation sequencing data. *BMC Bioinf.* 11, 485 (2010).
- E. Kopylova, L. Noé, H. Touzet, SortMeRNA: Fast and accurate filtering of ribosomal RNAs in metatranscriptomic data. *Bioinformatics* 28, 3211–3217 (2012).
- X. Su, W. Pan, B. Song, J. Xu, K. Ning, Parallel-META 2.0: Enhanced metagenomic data analysis with functional annotation, high performance computing and advanced visualization. *PLoS One* 9, e89323 (2014).
- C. Quast et al., The SILVA ribosomal RNA gene database project: Improved data processing and web-based tools. Nucleic Acids Res. 41, D590–D596 (2013).
- 64. C. Camacho et al., BLAST+: Architecture and applications. BMC Bioinf. 10, 421 (2009).
- J. G. Caporaso et al., QIIME allows analysis of high-throughput community sequencing data. Nat. Methods 7, 335–336 (2010).
- R. C. Edgar, Search and clustering orders of magnitude faster than BLAST. *Bio-informatics* 26, 2460–2461 (2010).
- M. N. Price, P. S. Dehal, A. P. Arkin, FastTree 2—Approximately maximum-likelihood trees for large alignments. *PLoS One* 5, e9490 (2010).
- UniProt Consortium, UniProt: A hub for protein information. Nucleic Acids Res. 43, D204–D212 (2015).
- B. L. Cantarel et al., The carbohydrate-active enZymes database (CAZy): An expert resource for glycogenomics. Nucleic Acids Res. 37, D233–D238 (2009).
- B. Buchfink, C. Xie, D. H. Huson, Fast and sensitive protein alignment using DI-AMOND. Nat. Methods 12, 59–60 (2015).
- Y. Peng, H. C. M. Leung, S. M. Yiu, F. Y. L. Chin, IDBA-UD: A de novo assembler for single-cell and metagenomic sequencing data with highly uneven depth. *Bio*informatics 28, 1420–1428 (2012).
- D. D. Kang, J. Froula, R. Egan, Z. Wang, MetaBAT, an efficient tool for accurately reconstructing single genomes from complex microbial communities. *PeerJ* 3, e1165 (2015).
- Y.-W. Wu, B. A. Simmons, S. W. Singer, MaxBin 2.0: An automated binning algorithm to recover genomes from multiple metagenomic datasets. *Bioinformatics* 32, 605–607 (2016).
- B. Langmead, S. L. Salzberg, Fast gapped-read alignment with Bowtie 2. Nat. Methods 9, 357–359 (2012).
- H. Li et al.; 1000 Genome Project Data Processing Subgroup, The sequence alignment/ map format and SAMtools. Bioinformatics 25, 2078–2079 (2009).
- D. Hyatt et al., Prodigal: Prokaryotic gene recognition and translation initiation site identification. BMC Bioinf. 11, 119 (2010).
- L. M. Rodriguez-R et al., The Microbial Genomes Atlas (MiGA) webserver: Taxonomic and gene diversity analysis of Archaea and Bacteria at the whole genome level. *Nucleic Acids Res.* 46, W282–W288 (2018).
- D. H. Parks et al., A standardized bacterial taxonomy based on genome phylogeny substantially revises the tree of life. Nat. Biotechnol. 36, 996–1004 (2018).
- J. Pinheiro, D. Bates, S. DebRoy, D. Sarkar; R Core Team, nlme: Linear and nonlinear mixed effects models (2018). https://cran.r-project.org/web/packages/nlme/index.html. Accessed 15 June 2018.
- M. I. Love, W. Huber, S. Anders, Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biol.* 15, 550 (2014).
- Y. Benjamini, Y. Hochberg, Controlling the false discovery rate: A practical and powerful approach to multiple testing. J. R. Stat. Soc. B 57, 289–300 (1995).
- R. L. Barter, B. Yu, Superheat: An R package for creating beautiful and extendable heatmaps for visualizing complex data. J. Comput. Graph. Stat. 27, 910–922 (2018).

ENVIRONMENTAL SCIENCES

Johnston et al.